Causes and consequences of polar amplification elucidated by coordinated multimodel experiments

James A. Screen¹, Alexandre Audette², Russell Blackport³, Clara Deser⁴, Mark England¹, Nicole Feldl², Melissa Gervais⁵, Stephanie Hay¹, Paul J. Kushner⁶, Yu-Chiao Liang⁷, Rym Msadek⁸, Regan Mudhar¹, Michael Sigmond³, Doug Smith⁹, Lantao Sun¹⁰ and Hao Yu¹

¹Department of Mathematics and Statistics, University of Exeter, Exeter, UK

²Department of Earth and Planetary Sciences, University of California Santa Cruz, Santa Cruz, USA

³Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, Canada.

⁴National Center for Atmospheric Research, Boulder, USA

⁵Department of Meteorology and Atmospheric Science, Penn State University, University Park, USA

⁶Department of Physics, University of Toronto, Toronto, Canada

⁷Department of Atmospheric Sciences, National Taiwan University, Taipei, Taiwan

⁸CECI, CNRS, IRD, CERFACS, Université de Toulouse, Toulouse, France

⁹Met Office Hadley Centre, Met Office, Exeter, UK

¹⁰Department of Atmospheric Science, Colorado State University, Fort Collins, USA

Abstract

Human-induced warming is amplified in the polar regions because of energy budget changes and feedbacks triggered by diminishing sea ice and snow cover. The Arctic has warmed three to four times faster than the global average in recent decades and is projected to continue to warm disproportionately. Arctic amplification can lead to changes in climate and weather patterns in the northern midlatitudes, but these remote consequences are not precisely known, due to difficulties determining cause and effect, separating forced climate change from internal climate variability, and discrepancies between studies. To address these challenges, the Polar Amplification Model Intercomparison Project (PAMIP) delivered a large set of coordinated multi-model experiments and analyses. Here, we review the main scientific advances facilitated by the PAMIP. Tropospheric responses to Arctic sea-ice loss that are robust across models and separable from internal variability have been revealed, including local warming and moistening, equatorward shifts of the jet stream and storm track in the North Atlantic, reduced atmospheric poleward heat transport, and fewer and milder cold extremes over North America. Whilst generally small compared to simulated internal variability, the response to future Arctic sea-ice loss comprises a non-negligible, and for some regions and variables, large, contribution to projected climate change. For example, Arctic sea-ice loss is essential to explain projected North Atlantic jet trends and their uncertainty. Model diversity in the simulated responses has provided pathways to observationally constrain the real-world response. Through novel analyses of existing experiments, new models and simulations, and enhanced collaboration, there are opportunities to gain further insights into the causes and consequences of polar amplification.

Introduction

Polar amplification describes the phenomenon that the polar regions warm at a faster rate than the global average in response to increased greenhouse gases¹⁻³. It has long been attributed to the surface albedo feedback: diminishing snow and sea ice reflects less incoming solar radiation, promoting warming and further snow and sea-ice loss. This intuitive explanation, however, is not the full picture for a few fundamental reasons. Firstly, polar amplification occurs in climate models even without any loss of ice or snow, albeit with reduced magnitude⁴. Second, feedback analysis of climate model output consistently highlights the lapse rate feedback, which characterises the sensitivity to the vertical temperature structure, as the single most important driver of polar amplification². Lastly, while sea-ice loss is greatest in late summer, polar amplification is most pronounced in winter⁵. We now understand that polar amplification is a coupled atmosphere-ocean-ice phenomenon that operates across the seasonal cycle^{6,7}. It is important to note that the framework through which polar amplification is viewed

can lead to different conclusions about the importance of different processes, as the relevant feedbacks are interconnected, but interact in complex ways, with the impact of individual feedbacks potentially enhanced through synergistic interactions⁸⁻¹⁰. Thus, although the processes and feedbacks leading to polar amplification are reasonably well understood, their relative contributions and how they lead to hemispheric and seasonal asymmetries are not precisely known^{6,7}.

Polar amplification may trigger remote climate responses in other parts of the world. For example, a direct consequence of Arctic amplification is a reduction in the near-surface meridional temperature gradient at high latitudes, which implies a weaker jet stream through thermal wind balance¹¹. A weaker jet stream may in turn affect the propagation of weather disturbances across the Northern Hemisphere midlatitudes, impacting regional climate¹²⁻¹⁴. However, there is a lack of scientific consensus on the details of any potential influence on midlatitudes^{11,13}. Major challenges here are to separate cause from effect, and forced changes from internal variability^{15,16}.

Climate models are useful tools to probe questions of causality and physical mechanisms, as they allow for controlled experiments to isolate and quantify specific forcings or processes. For example, an atmospheric model can be provided with different sea ice conditions to isolate the atmospheric response to sea-ice loss. Myriad sea ice perturbation experiments have been conducted, but often with apparently conflicting results^{12,13,17}. For example, models seemingly disagree on the sign of regional winter temperature changes over midlatitudes in response to sea-ice loss¹⁷⁻¹⁹. Some of the apparent differences between models could relate to sampling uncertainty or due to varying aspects of the experimental design. However, even absent these factors, the forced response may still be model dependent. Understanding if and why models differ is important to constrain projections of future climate change²⁰.

To address these questions, the Polar Amplification Model Intercomparison Project (PAMIP) provided a framework for the scientific community to produce and analyse coordinated model experiments²⁰. The PAMIP experiments broadly fall into two categories (**Fig. 1**): experiments with perturbed sea ice cover (prescribed in atmosphere-only experiments and nudged in coupled experiments) but baseline sea surface temperatures (SST), and experiments with perturbed SST but baseline sea ice. Differencing these experiments allows for the quantification of the simulated responses to sea ice, separately in the Arctic and Antarctic, or SST change, absent other factors (Fig. 1). The 'Tier 1' time-slice simulations have been most widely run and utilised. These are year-long simulations with prescribed SST and sea ice representing preindustrial, present, or future time periods, each with at least 100 ensemble members (i.e., repeated simulations), where ensemble members differ only by very small changes in their initial conditions. Such a large sample size ensures a more robust quantification of the simulated response to sea ice and SST change in the face of internal variability. This, in turn, allows for a comparison of the forced response across models to quantify model uncertainty and, through the identification of emergent relationships across models and the application of observational constraints, to potentially narrow uncertainties in the real-world response.

The purpose of this Perspective is to synthesise recent advances in understanding the causes and consequences of polar amplification facilitated by the PAMIP. It is not our intention to provide a comprehensive review of polar amplification, which can be found elsewhere^{6,7,12,13}. Instead, we draw upon and summarise the latest developments in the field (focussing on work published since the inception of PAMIP in 2019) to identify key advances, remaining questions and pathways to address these, including the need for new coordinated model experiments.

Causes of Arctic amplification

The PAMIP experiments serve as an ideal testbed to further understand the processes responsible for Arctic amplification. Jenkins et al. (2024) used the PAMIP experiments to show the roles of Arctic seaice loss and global SST change in contributing to different local feedbacks and remote processes²¹. Arctic warming in response to sea-ice loss maximises in winter, due to greatly enhanced oceanic heat release. This produces lower tropospheric Arctic warming and triggers positive lapse rate and cloud feedbacks, leading to large Arctic amplification (**Fig. 2**). Despite strong albedo feedback in summer, atmospheric warming is muted. In contrast, in response to global SST warming absent sea-ice loss, enhanced atmospheric energy convergence into the Arctic is the dominant contributor to Arctic warming, although Arctic amplification is relatively small compared to that in response to sea-ice loss (**Fig. 2**). As SST-induced warming is larger aloft than at the surface, the lapse rate feedback is negative for much of the year. The water vapor feedback contributes to Arctic warming but opposes amplification due to larger tropical than Arctic moistening under SST-induced warming with fixed Arctic sea-ice. These results reinforce that Arctic amplification is caused primarily by sea-ice loss and resultant local changes in surface fluxes, while increased poleward energy transport can only produce weak amplification in the absence of sea-ice-related feedbacks.

While the magnitude and seasonality of polar amplification are controlled by local processes and feedbacks (**Fig. 2**), disproportionately large Arctic warming is a fundamental response of a moist atmosphere to an increase in greenhouse gases^{22,23}. A preferential increase in tropical humidity, which occurs due to the Clausius-Clapeyron relation, provokes an increase in poleward atmospheric latent energy transport. This process explains polar amplification in the absence of sea ice and related polar feedbacks^{4,24}. PAMIP has expanded understanding of these poleward energy transport changes by linking them to the processes of sea-ice loss and SST changes. Audette et al. (2021) shows that Arctic sea-ice loss reduces northward eddy-driven energy transport into the Arctic in all the PAMIP models, owing to the reduced near-surface temperature gradient, that is balanced by a similar magnitude increase

of latent energy transport due to SST warming²⁵. This opposition in total energy transport mainly arises in the annual mean as the influence of Arctic sea-ice loss is greatest in winter while that of SST warming is greatest in summer²⁶. Hence, there is growing appreciation that different aspects of the poleward energy transport exhibit different efficacies, and that the polar-cap-averaged energetic perspective risks obscuring the role of remote processes in driving polar amplification.

Precipitation changes are also amplified in the Arctic relative to lower latitudes²⁷. The PAMIP experiments have shown that sea-ice loss and SST warming both increase Arctic precipitation, but through predominantly different mechanisms. SST warming increases precipitation in the Arctic²⁸, consistent with the increase in moisture convergence in the same experiments^{25,26}, while sea-ice loss also increases precipitation through increased evaporation^{21,28,29}. Sea-ice loss is critical for amplified wetting in the Arctic, as SST warming alone leads to larger wetting at lower than higher latitudes²⁹.

Robust remote responses to sea-ice loss

Prior to the PAMIP, the precise nature of the remote circulation response to sea ice loss, including the effect on the jetstream, storm tracks and dominant modes of variability (e.g., North Atlantic Oscillation, Northern Annular Mode), as well as the mechanisms involved were elusive^{13,20}. This is due to a variety of factors, including, but not limited to, inter-model differences, differences in experimental design, and sampling errors due to large internal variability. The PAMIP has facilitated like-for-like comparison across models (owing to identical boundary conditions) and better separation of the forced response from internal variability (by requiring a minimum of 100 ensemble members), although even larger ensembles appear to be necessary to confidently capture the stratospheric response³⁰⁻³² and changes in regional weather extremes³³⁻³⁵, as discussed later. One important advance from the PAMIP has been to identify robust large-scale troposphere circulation responses to future Arctic³⁶ and Antarctic sea-ice loss³⁷.

Smith et al. (2022) examined the PAMIP 'present-day' baseline and future Arctic sea ice experiments from 16 atmospheric models and showed a robust equatorward shift of the tropospheric westerly jet in response to future Arctic sea-ice loss, albeit of varying magnitudes between models³⁶. A similar equatorward shift of the wintertime jet was found in the southern hemisphere in response to future Antarctic sea-ice loss, also robust in sign but of varying magnitude across models³⁷. The mechanisms that lead to an equatorward jet shift are shown schematically in **Fig. 3**. Briefly, Arctic warming induced by sea-ice loss reduces the high-latitude meridional temperature gradient (step 1 in **Fig. 3**), reducing wind speed on the poleward flank of the jet (step 2; following the thermal wind relationship). The weakened temperature gradient also reduces baroclinic eddy activity, weakening the storm track and in turn, reducing the upward wave activity flux from the surface (consistent with reduced eddy-driven

poleward heat transport). An anomalous meridional circulation develops with ascent around 40-50°N, poleward flow in the mid to upper troposphere, descent around 65–75°N, and equatorward flow near the surface (step 3). Adiabatic cooling of the ascending branch acts to enhance the latitudinal temperature gradient on the equatorward side of the jet, which strengthens the wind to the south of the jet core (step 4). Taken together, the weakened westerlies to the north and strengthened westerlies to the south imply an equatorward shift of the jet and storm activity (step 5). The shifted storm activity leads to wave flux anomalies that reinforce the jet shift, through positive eddy feedback (step 6). The magnitude of the response across models appears to depend on the strength of this atmospheric eddy feedback, in both hemispheres^{36,37}. This zonal-mean perspective applies also for the Atlantic basin. In the Pacific, while the anomalies are broadly similar, the climatological jet is located farther south, so the westerly wind anomalies act to strengthen the Pacific jet rather than shift it equatorward³⁸.

Regionally, the winter North Atlantic jet shifts equatorward in response to Arctic sea-ice loss in most models, with a few models showing a negligible response^{35,30,40}. However, models disagree on the sign of simulated changes in the speed and tilt of the North Atlantic jet with most models showing negligible change^{35,39,40}. In contrast, the North Pacific jet is simulated to strengthen in response to future Arctic sea-ice loss, on average across models^{38,41}. The contrasting jet speed responses in the North Atlantic and North Pacific can be understood by the more equatorward climatological position of the North Pacific jet, such that the strengthened westerly wind in response to sea-ice loss occurs in the core of the jet in the Pacific rather than on the equatorward flank, as simulated in the Atlantic^{38,42}.

Accompanying the jet shift is a weakening and equatorward shift of the North Atlantic storm track, revealed by both Lagrangian (storm identification and tracking) and Eulerian (grid-point sea level pressure variability) approaches^{34,38,43,44}. An eastward shift or extension of the North Pacific storm track is consistent with a strengthened and extended jet^{38,41}. Across the whole Northern Hemisphere, in response to future Arctic sea-ice loss there are fewer individual storms simulated, those storms are weaker, they propagate more slowly, and have longer lifetimes⁴³, which can be understood as an energetic response to change in surface albedo and weakened poleward atmospheric energy transport⁴⁵. However, over North America, weather systems do not appear to stagnate, although they tend to be weaker³³. Other weather system responses to Arctic sea-ice loss identified using the PAMIP experiments include an increase in winter Scandinavian blocking^{35,46}, a reduction in cut-off-lows over Southern Europe⁴⁷, enhanced tropical cyclone genesis over the eastern North Pacific⁴⁸, and shifts in the predominant locations of atmospheric rivers⁴⁹.

Drying over Northwest Europe and wetting over central Europe are found in response to future Arctic sea-ice loss, closely linked to storm track changes^{34,35,39,50}. Warmer and fewer cold air outbreaks over central and western North America are connected to the strengthened and extended North Pacific jet⁴¹.

Reduced meridional winds are linked to increased surface temperature persistence⁵¹. In addition to these dynamically driven responses, advection across the weakened meridional temperature gradient causes a reduction in day-to-day temperature variability and more rapid warming of cold extremes than of the average temperature⁵², consistent with earlier findings⁵³. Cold days, which typically occur under northerly flow, warm faster than the seasonal mean (and hot days), as northerly flow advects Arctic air masses that are significantly warmed by sea-ice loss to lower latitudes⁵². Lo et al. (2023) found the 1-in-20-year extreme cold events over northern mid-to-high latitudes warmed by up to 2.5°C for eastern Canada and the northeast United States⁵⁴.

The varied and widespread responses to Arctic sea-ice loss, summarised visually in **Fig. 4**, have been predominantly identified in winter, although some responses have been reported in summer^{48,55-58}. The larger magnitude responses in winter than summer^{9,28,43}, arise due to the seasonality of the Arctic surface energy budget response to sea-ice loss^{9,59}. Anomalous energy transfer from the ocean to atmosphere maximises in the winter, as does the Arctic warming response, which is the trigger for the subsequent effects on the atmospheric circulation and midlatitude weather and climate.

Reducing and exploiting uncertainty

Atmospheric internal variability has emerged as a common challenge in understanding the character of the forced response to sea-ice loss, especially for applications involving regional climate and extremes^{33-35,41,49,60} and the stratospheric polar vortex^{30-32,61,62}. The PAMIP protocol recommended a minimum ensemble size of 100 members. However, Peings et al. (2021) found substantial inconsistencies in the atmospheric circulation response to Arctic sea-ice loss among three separate 100member ensembles, indicating that ensembles of this size still contain a significant imprint of internal variability³⁰. Similarly, Sun et al. (2022) showed that the stratospheric polar vortex response in PAMIP simulations is small compared to its internal variability³¹. Even the sign of the stratospheric circulation response to sea-ice loss is uncertain³⁶, for reasons that remain unclear but are likely to include low signal-to-noise and state dependence^{32,62}, and in turn influences the magnitude of the tropospheric circulation response^{31,32,62}. Ye et al. (2024) is unique in conducting very large ensembles (~2000 members) with two models following the PAMIP protocol and found that several hundred members (> 400) are needed to robustly estimate the seasonal-mean large-scale circulation response, and a thousand or more members for regional climate extremes³⁴. This could imply the real-world response to sea-ice loss is weak^{15,36,37,63}. However, the weak simulated responses in the PAMIP experiments could be a symptom of a broader signal-to-noise problem in models⁶⁴ and thus, the real-world response may be larger than models suggest.

Internal variability appears insufficient to fully explain the model spread in the magnitude of the tropospheric circulation response^{36,37}, suggesting this response is model dependent. One factor that may lead to model dependence is the strength of atmospheric eddy feedback: models exhibiting stronger eddy feedback simulate a stronger tropospheric circulation response to Arctic and Antarctic sea-ice loss^{36,37}. Understanding the causes of model spread is valuable not only for interpreting model responses but also for reducing uncertainty in projections through an emergent constraint—an inter-model relationship linking an observable aspect of the climate to the response to sea-ice loss across the PAMIP models^{36,37}. The observed estimate of eddy feedback is larger than any of the modelled values, which suggests that models systematically underestimate the atmospheric circulation response to Arctic sea-ice loss. While there are suggestions that simulated eddy feedback is too weak owing to coarse model resolution⁶⁵, it remains unclear if the response to sea-ice loss is dependent on model resolution⁶¹. Constraining by the observed eddy feedback increases the equatorward jet shift in response to Arctic sea-ice loss by 40% compared to the multimodel mean, becoming of larger magnitude than the simulated poleward shift in response to SST warming³⁷.

Additionally, differences in models' unperturbed climate - often referred to as the basic state - can also influence the response to sea-ice loss^{62,66}. The basic state of the winds in the so-called "neck region", located in the upper troposphere-lower stratosphere between the latitudes of the subtropical jet and polar jet, determines whether the stratospheric pathway is active, and thereby affects the magnitude of the tropospheric circulation response to Arctic sea-ice loss^{62,67}. Work is underway to provide an alternative emergent constraint based on the strength of the neck region winds⁶². Stratosphere-troposphere coupling also plays a central role in mid-tropospheric Arctic warming in response to sea-ice loss⁶⁸, and this deep warming, can further enhance the tropospheric circulation response⁶⁹.

Slow modes of natural climate variability can alter the 'basic state' and thereby, modulate the response to sea-ice loss. Labe et al. (2019) found that the atmospheric circulation response to Arctic sea-ice loss was stronger during the easterly phase of the Quasi-Biennial Oscillation (QBO) than its westerly phase⁷⁰. This led one group, the UK Met Office, to perform separate PAMIP ensembles for different QBO phases. The stratospheric polar vortex weakened in response to Arctic sea-ice loss during easterly QBO but not during westerly QBO^{32,71,72}. The different QBO states across the PAMIP models may contribute to the model differences in the responses of the stratospheric polar vortex³². There is also evidence that the phase of the El Niño-Southern Oscillation (ENSO), Pacific (Inter-) Decadal Variability (PDV), and Atlantic Multidecadal Variability (AMV) all influence the atmospheric response to Arctic sea-ice loss^{30,72-76}. These modes of variability rely on coupling between the ocean and atmosphere. A recent study by Cvijanovic et al. (2025) suggests that the North Pacific response to Arctic sea-ice loss is strongly influenced by ocean–atmosphere coupling in the tropics⁷⁷. Models that prescribe

tropical SSTs — thereby lacking ocean-atmosphere coupling - tend to simulate a strengthened Aleutian Low over the North Pacific in response to sea-ice loss³⁶, as do many coupled model experiments¹⁷, but in the study of Cvijanovic et al. (2025), the inclusion of coupling resulted in weakened Aleutian Low⁷⁷. We will return to the potential importance of ocean coupling and the limitations of prescribing climatological SST (i.e. lacking internal ocean variability) later.

Interpreting climate projections

Confidence in climate projections is increased when the underpinning physical mechanisms are well understood. Since sea-ice loss and SST warming may affect the atmospheric circulation in different ways and through different mechanisms, it is of value therefore, to consider the responses to these factors separately, which the PAMIP experiments has facilitated. The effect of SST warming dominates over most of the globe, outside the Arctic^{9,28} (**Fig. 5**). However, the effect of sea-ice loss is as important, if not more, than SST warming for response of the winter North Atlantic circulation^{28,35,38} (**Fig. 5**) and East Asian summer monsoon⁵⁸.

Separation of the responses to SST warming and sea-ice loss has been particularly useful to uncover the mechanisms driving the extratropical circulation changes. Using the PAMIP experiments, Yu et al. (2024) demonstrated a significant weakening of the westerlies over the high-latitude North Atlantic and strengthening over the midlatitude North Atlantic in response to Arctic sea-ice loss, while ocean warming led to a broadly opposite response²⁸ (**Fig. 5**). This regional 'tug-of-war' in the Atlantic is also seen for storm track activity^{38,50}, but not in the Pacific^{28,38,41}. Hay et al (2025) showed that more equatorward average location of the Pacific jet and the geography of the North Pacific results in circulation responses to sea-ice loss and ocean warming that reinforce rather than oppose each other³⁸.

The role of Arctic sea-ice loss on cold winter extremes, in the context of a warming climate, has been the subject of debate for the past few decades¹²⁻¹⁵. The PAMIP has facilitated a clearer picture on the relative roles of SST warming and Arctic sea-ice loss on mid-latitude extreme cold events. While Arctic sea-ice loss leads to more frequent or severe cold extremes in some simulations in limited midlatitude regions^{19,54}, these changes are small and overwhelmed by strong warming of extreme cold events over the midlatitudes in response to ocean warming^{46,54}. However, sea-ice loss plays a more dominant role than SST warming in the reduction of temperature variability over North America⁵².

It is worth noting that the PAMIP experiments are based upon one scenario of sea-ice loss and SST warming, but model projections suggest a range of outcomes. In the PAMIP experiments, comparing the annual-mean future and present-day states, there is a loss of 3.0 million km² of sea ice cover for a global SST warming of 1.4 C, equating to approximately 2.1 million km² of sea-ice loss per degree

Celsius of global SST warming. Model projections show sea-ice sensitivities of less than 1 to more than 4 million km² per degree of global SST warming³⁵. Under the assumptions of linear additivity and scalability²⁹, the PAMIP-derived responses to sea-ice loss and SST warming can be scaled and summed to illustrate different storylines, with lesser or greater sea-ice sensitivities (**Fig. 6**), which is closely related to model uncertainty in the magnitude of Arctic amplification. Arctic warming, Arctic amplification (2.7 and 4.2 in low and high sensitivity cases, respectively) and Arctic wetting (0.12 and 0.18 mm/day, respectively) increase with greater sea-ice sensitivity to global warming (**Fig. 6**). The magnitude and pattern of the circulation response at lower latitudes are also sensitive to the storyline. For example, the North Atlantic jet shifts marginally poleward (0.1° latitude) in the low sea-ice sensitivity storyline and equatorward (-0.5°) in the high sea-ice sensitivity storyline. Given that models appear to underestimate the observed sea-ice sensitivity⁷⁸, the high sea-ice sensitivity storyline may be the most plausible.

Future directions

To close, we highlight some future research opportunities.

Hemispheric asymmetry. First, polar amplification exhibits notable hemispheric asymmetry, with the Arctic experiencing significantly stronger warming than the Antarctic². Several mechanisms underpinning this disparity have been proposed, including a buffering effect by the presence of a deep and cold Southern Ocean, which limits surface warming through effective oceanic heat uptake79, and high topography in the Antarctic that inhibits temperature longwave feedbacks¹. At a longer timescale, oceanic heat transport contributes to this hemispheric asymmetry, as the AMOC facilitates poleward heat transfer and local feedbacks in the Northern Hemisphere⁸⁰, whereas analogous mechanisms are weaker in the Southern Hemisphere. Beyond temperature change, Arctic precipitation amplification driven by increased moisture availability and enhanced poleward moisture transport that remains energetically constrained^{27,81} - enhances Arctic hydrological sensitivity and serves as another manifestation of polar amplified climate change. In sum, the interplay between sea-ice feedbacks, ocean dynamics, and atmospheric processes leads to pronounced and hemispherically asymmetric polar climate responses. We urge more process-based analysis to improve our understanding of asymmetries in the two poles and their implications for global climate. This need is particularly urgent given the unprecedented Antarctic sea-ice loss since 2016, which may signal a transition to a new Antarctic climate regime⁸²⁻⁸⁴, for which the associated atmospheric response, both locally and globally, remains relatively unknown^{85,86}. Advancing our understanding of these processes is critical for anticipating future climate impacts.

Experimental protocols. Within the framework of the PAMIP, a variety of experimental strategies have been employed to advance the understanding of polar climate processes and their remote influences, including prescribing ocean surface boundary conditions (in atmosphere-only experiments) and nudging or albedo reduction techniques (in coupled atmosphere-ocean-ice experiments). Ocean coupling appears to modulate the response to sea-ice loss and aspects of the response may be underestimated by prescribing sea ice and thereby inhibiting ocean-ice-atmosphere interaction^{42,44,85,87-}

⁹⁰. The PAMIP coupled experiments simulate enhanced weakening of the winter storm tracks in response to sea-ice loss, compared to the uncoupled experiments⁴⁴, and deeper Arctic warming and enhanced strengthening of the Siberian High⁹⁰. However, concerns have been raised that the methods used to constrain sea ice in coupled modes might lead to spurious responses⁹¹⁻⁹⁴. New methods for addressing these issues are necessary for developing protocols for the next phase of the PAMIP. For example, the climate impacts of spurious heating associated with current nudging techniques can be quantified, and hence removed, through additional model simulations⁹² or through post processing via pattern scaling⁹⁴. Alternatively, regional CO₂ forcing may offer a pathway for isolating Arctic change with fewer unintended consequences and is technically more straightforward to implement across all climate models than nudging^{95,96}. Other methods, such as those involving modifications to sea ice and snow-over-sea-ice parameters^{77,84,97} or atmosphere-ocean-ice coupling⁹⁸, have been proposed and may reduce any spurious responses. The most utilised experiments, the year-long atmosphere-only simulations, not only omit ocean coupling, but also lack ocean internal variability (as climatological boundary conditions are prescribed) and processes acting on interannual timescales. Transient experiments, although more costly to run, allow for better sampling of internal variability and response timescales. Additionally, considering different initial states in the presence of sea ice forcings, such sampling opposite phases of the QBO, ENSO, AMV or PDV, would further enhance our understanding of the state dependency (i.e. nonlinearity) of responses to sea-ice loss⁷⁰⁻⁷⁶. Greater coordination of atmospheric and ocean initial conditions across modelling protocols may help to explain apparent model discrepancies that remain hard to reconcile, for example, the divergent stratospheric polar vortex responses across models³².

New models and experiments. High-resolution models, either globally or with regional grid refinement, better capture fine-scale processes and therefore further improve the representation of sea-ice-atmosphere interactions and feedbacks^{29,99} and simulate stronger circulation responses to SST anomalies¹⁰⁰. However, whether the atmospheric response to sea-ice loss is sensitive to model resolution remains unclear^{29,32,61} and warrants further investigation. Additionally, saving more diagnostic variables at high temporal frequency (daily or 6-hourly), would provide opportunities to examine processes at the scale of weather events. New experiments with different prescribed boundary conditions (e.g., larger global warming and sea-ice loss to increase the signal-to-noise ratio, and different storylines of sea ice and SST change) may also yield important insights. Repeating the existing experiments with newer

models, including at higher resolution, would be valuable to test the robustness of emergent constraints with independent ensembles. Idealized simulations also continue to serve as a useful tool for conceptual understanding, providing controlled environments to disentangle nonlinear interactions and benchmark model behaviour across scales^{1,4,21,24,45,51,67,93,101}. More recently, Artificial Intelligence and machine learning-based climate models¹⁰² offer new avenues to simulate complex climate dynamics with enhanced computational efficiency, although their capacity to fully capture sea-ice thermodynamics and large-scale coupling remains under assessment.

Enhanced collaboration. We encourage the use of PAMIP simulations for novel and varied applications. Exemplars include interpreting the implications of incomplete Arctic sea-ice recovery under CO₂ removal¹⁰³, exploring the role of sea-ice-related feedbacks in sustaining Barents Sea ice loss¹⁰⁴, identifying the surface signature of stratospheric variability¹⁰⁵, and quantifying the effects of projected sea-ice loss on cold-related mortality¹⁰⁶, Greenland ice sheet mass balance⁵⁷, Yangtze River Basin heatwaves¹⁰⁷ and eastern Siberian wildfires¹⁰⁸, amongst others. Expanding collaborations with other research communities will be critical for the next phase of PAMIP. Coordination between future PAMIP and other model intercomparison projects could enhance process-based understanding. Joint efforts with the climate feedback, atmosphere-ocean-ice dynamics, and polar process communities could further enrich the interpretation of results. Side-by-side analysis of modelling experiments within dynamical and radiative feedback frameworks will provide deeper insights into the processes and mechanisms by which a climate response to sea ice loss is provoked. Closer ties with the polar observational community would support robust model–observation comparisons, which likely needs better experimental designs and additional high-frequency model outputs.

In summary, the PAMIP has facilitated a plethora of studies (not all are cited here), but arguably the biggest collective advance has been to reveal robust tropospheric responses to Arctic sea-ice loss, that whilst generally small compared to simulated internal variability, comprise a non-negligible, and for some regions and variables, large (relative to that of SST warming), contribution to projected climate change. Further, it has uncovered and explained responses that are dependent on model physics and/or background states, which provide pathways to constrain the real-world response. Looking forward, there are opportunities, through novel analyses of existing experiments, new simulations and model versions, and enhanced collaborations, to gain further insights into the causes and consequences of polar amplification.

Acknowledgements

Matt Jenkins is thanked for providing Fig. 2. We thank the modelling groups that have contributed to the PAMIP, and the Earth System Grid Federation (https://esgf.github.io) for providing the infrastructure for data storage and sharing. This work used JASMIN, the UK's collaborative data

analysis environment (https://www.jasmin.ac.uk). J.A.S and S.H were supported by the NERC ArctiCONNECT project (NE/V005855/1). C.D. was supported by the National Center for Atmospheric Research (NCAR), which is sponsored by the National Science Foundation under Cooperative Agreement 1852977. M.E was supported by the Royal Commission for the Exhibition of 1851 research fellowship. N.F. was supported by NSF Grant AGS-1753034. P.K. acknowledges support from the grant NSF PHY-2309135 to the Kavli Institute for Theoretical Physics (KITP). Y.-C. L. was supported by the National Science and Technology Council (113-2628-M-002-018 and 113-2116-M-008-024). R.M. is funded by a NERC GW4+ Doctoral Training Partnership studentship (NE/S007504/1). D.S. was supported by the Met Office Hadley Centre Climate Programme funded by DSIT. L.S. was supported by NSF Grant AGS-2300038. H.Y. was supported by the Chinese Scholarship Council.

References

- Hahn, L. C., Armour, K. C., Battisti, D. S., Donohoe, A., Pauling, A. G. & Bitz, C. M. Antarctic Elevation Drives Hemispheric Asymmetry in Polar Lapse Rate Climatology and Feedback. Geophys. Res. Lett. 47, (2020), <u>https://doi.org/10.1029/2020GL088965</u>
- Hahn, L. C., Armour, K. C., Zelinka, M. D., Bitz, C. M. & Donohoe, A. Contributions to Polar Amplification in CMIP5 and CMIP6 Models. Front. Earth Sci. 9, (2021), <u>https://doi.org/10.3389/feart.2021.710036</u>
- Rantanen, M., Karpechko, A. Yu., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T. & Laaksonen, A. The Arctic has warmed nearly four times faster than the globe since 1979. Commun. Earth Environ. 3, (2022), <u>https://doi.org/10.1038/s43247-022-00498-3</u>
- England, M. R. & Feldl, N. Robust Polar Amplification in Ice-Free Climates Relies on Ocean Heat Transport and Cloud Radiative Effects. J. Clim. 37, 2179–2197 (2024), <u>https://doi.org/10.1175/JCLI-D-23-0151.1</u>
- Screen, J. A. & Simmonds, I. The central role of diminishing sea ice in recent Arctic temperature amplification. Nature 464, 1334–1337 (2010), https://doi.org/10.1038/nature09051
- Previdi, M., Smith, K. L. & Polvani, L. M. Arctic amplification of climate change: a review of underlying mechanisms. Environ. Res. Lett. 16, 093003 (2021), <u>https://doi.org/10.1088/1748-9326/ac1c29</u>
- Taylor, P. C., Boeke, R. C., Boisvert, L. N., Feldl, N., Henry, M., Huang, Y., Langen, P. L., Liu, W., Pithan, F., Sejas, S. A. & Tan, I. Process Drivers, Inter-Model Spread, and the Path Forward: A Review of Amplified Arctic Warming. Front. Earth Sci. 9, (2022), <u>https://doi.org/10.3389/feart.2021.758361</u>

- Feldl, N., Po-Chedley, S., Singh, H. K. A., Hay, S. & Kushner, P. J. Sea ice and atmospheric circulation shape the high-latitude lapse rate feedback. npj Clim. Atmos. Sci. 3, (2020), https://doi.org/10.1038/s41612-020-00146-7
- Jenkins, M. & Dai, A. The Impact of Sea-Ice Loss on Arctic Climate Feedbacks and Their Role for Arctic Amplification. Geophys. Res. Lett. 48, (2021), https://doi.org/10.1029/2021GL094599
- Bonan, D. B., Kay, J. E., Feldl, N., & Zelinka, M. D. Mid-latitude clouds contribute to Arctic amplification via interactions with other climate feedbacks. Environ. Res. Clim. 4, 015001 (2025), <u>https://doi.org/10.1088/2752-5295/ada84b</u>
- Barnes, E. A. & Screen, J. A. The impact of Arctic warming on the midlatitude jet-stream: Can it? Has it? Will it? WIREs Clim. Change 6, 277–286 (2015), <u>https://doi.org/10.1002/wcc.337</u>
- Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., & Jones, J. Recent Arctic amplification and extreme mid-latitude weather. Nat. Geosci. 7, 627–637 (2014), https://doi.org/10.1038/ngeo2234
- Cohen, J., Zhang, X., Francis, J., Jung, T., Kwok, R., Overland, J., Ballinger, T. J., Bhatt, U. S., Chen, H. W., Coumou, D., Feldstein, S., Gu, H., Handorf, D., Henderson, G., Ionita, M., Kretschmer, M., Laliberte, F., Lee, S., Linderholm, H. W., Maslowski, W., Peings, Y., Pfeiffer, K., Rigor, I., Semmler, T., Stroeve, J., Taylor, P. C., Vavrus, S., Vihma, T., Wang, S., Wendisch, M., Wu, Y., & Yoon, J. Divergent consensuses on Arctic amplification influence on midlatitude severe winter weather. Nature Clim. Change 10, 20–29 (2020), https://doi.org/10.1038/s41558-019-0662-y
- Hanna, E., Francis, J., Wang, M., Overland, J. E., Cohen, J., Luo, D., Vihma, T., Fu, Q., Hall, R. J., Jaiser, R., Kim, S.-J., Köhler, R., Luu, L., Shen, X., Erner, I., Ukita, J., Yao, Y., Ye, K., Choi, H. & Skific, N. Influence of high-latitude blocking and the northern stratospheric polar vortex on cold-air outbreaks under Arctic amplification of global warming. Environ. Res.: Clim. 3, 042004 (2024), https://doi.org/10.1088/2752-5295/ad93f3
- Blackport, R., Screen, J. A., van der Wiel, K., & Bintanja, R. Minimal influence of reduced Arctic sea ice on coincident cold winters in mid-latitudes. Nature Clim. Change 9, 697–704 (2019), <u>https://doi.org/10.1038/s41558-019-0551-4</u>
- Blackport, R. & Screen, J. A. Observed Statistical Connections Overestimate the Causal Effects of Arctic Sea Ice Changes on Midlatitude Winter Climate. J. Clim. 34, 3021–3038 (2021), <u>https://doi.org/10.1175/JCLI-D-20-0293.1</u>
- Screen, J. A., Deser, C., Smith, D. M., Zhang, X., Blackport, R., Kushner, P. J., Oudar, T., McCusker, K. E. & Sun, L. Consistency and discrepancy in the atmospheric response to

Arctic sea-ice loss across climate models. Nature Geosci. 11, 155–163 (2018), https://doi.org/10.1038/s41561-018-0059-y

- Screen, J. A. & Blackport, R. How Robust is the Atmospheric Response to Projected Arctic Sea Ice Loss Across Climate Models? Geophys. Res. Lett. 46, 11406–11415 (2019), https://doi.org/10.1029/2019GL084936
- Zheng, C., Wu, Y., Ting, M., Screen, J. A. & Zhang, P. Diverse Eurasian Temperature Responses to Arctic Sea Ice Loss in Models due to Varying Balance between Dynamic Cooling and Thermodynamic Warming. J. Clim. 36, 8347–8364 (2023), https://doi.org/10.1175/JCLI-D-22-0937.1
- Smith, D. M., Screen, J. A., Deser, C., Cohen, J., Fyfe, J. C., García-Serrano, J., Jung, T., Kattsov, V., Matei, D., Msadek, R., Peings, Y., Sigmond, M., Ukita, J., Yoon, J.-H. & Zhang, X. The Polar Amplification Model Intercomparison Project (PAMIP) contribution to CMIP6: investigating the causes and consequences of polar amplification. Geosci. Model Dev. 12, 1139–1164 (2019), <u>https://doi.org/10.5194/gmd-12-1139-2019</u>
- Jenkins, M. T., Dai, A. & Deser, C. Arctic climate feedback response to local sea-ice concentration and remote sea surface temperature changes in PAMIP simulations. Clim. Dyn. 62, 10599–10620 (2024), <u>https://doi.org/10.1007/s00382-024-07465-y</u>
- Feldl, N. & Merlis, T. M. Polar Amplification in Idealized Climates: The Role of Ice, Moisture, and Seasons. Geophys. Res. Lett. 48, (2021), https://doi.org/10.1029/2021GL094130
- Zhou, W., Leung, L. R., Xie, S.-P. & Lu, J. An analytic theory for the degree of Arctic Amplification. Nature Commun. 15, (2024), <u>https://doi.org/10.1038/s41467-024-48469-w</u>
- Henry, M., Merlis, T. M., Lutsko, N. J. & Rose, B. E. J. Decomposing the Drivers of Polar Amplification with a Single-Column Model. J. Clim. 34, 2355–2365 (2021), <u>https://doi.org/10.1175/JCLI-D-20-0178.1</u>
- Audette, A., Fajber, R. A., Kushner, P. J., Wu, Y., Peings, Y., Magnusdottir, G., Eade, R., Sigmond, M., & Sun, L. Opposite responses of the dry and moist eddy heat transport into the Arctic in the PAMIP experiments. Geophys. Res. Lett. 48, e2020GL089990 (2021), https://doi.org/10.1029/2020GL089990
- Hahn, L. C., Armour, K. C., Battisti, D. S., Donohoe, A. & Fajber, R. Seasonal Changes in Atmospheric Heat Transport to the Arctic Under Increased CO2. Geophys. Res. Lett. 50, (2023), <u>https://doi.org/10.1029/2023GL105156</u>
- McCrystall, M. R., Stroeve, J., Serreze, M., Forbes, B. C. & Screen, J. A. New climate models reveal faster and larger increases in Arctic precipitation than previously projected. Nature Commun. 12, (2021), <u>https://doi.org/10.1038/s41467-021-27031-y</u>
- 28. Yu, H., Screen, J. A., Xu, M., Hay, S. & Catto, J. L. Comparing the Atmospheric Responses to Reduced Arctic Sea Ice, a Warmer Ocean, and Increased CO2 and Their Contributions to

Projected Change at 2°C Global Warming. J. Clim.37, 6367–6380 (2024), https://doi.org/10.1175/JCLI-D-24-0104.1

- Sun, L., Wills, R. C. J, Deser. C et al. Increased model resolution amplifies Arctic precipitation and atmospheric circulation response to sea-ice loss. J. Clim (2025), submitted <u>https://doi.org/10.22541/essoar.175157568.87832579/v1</u>
- Peings, Y., Labe, Z. M. & Magnusdottir, G. Are 100 Ensemble Members Enough to Capture the Remote Atmospheric Response to +2°C Arctic Sea Ice Loss? J. Clim. 34, 3751–3769 (2021), <u>https://doi.org/10.1175/JCLI-D-20-0613.1</u>
- Sun, L., Deser, C., Simpson, I. & Sigmond, M. Uncertainty in the Winter Tropospheric Response to Arctic Sea Ice Loss: The Role of Stratospheric Polar Vortex Internal Variability. J. Clim. 35, 3109–3130 (2022), <u>https://doi.org/10.1175/JCLI-D-21-0543.1</u>
- Mudhar, R., Seviour, W. J. M., Screen, J. A., Thomson, S. I. & Turrell, C. Diverse and weak simulated stratospheric responses to future Arctic sea-ice loss. J. Geophys. Res. Atmos. (2025), submitted
- Gervais, M., Sun, L. & Deser, C. Impacts of Projected Arctic Sea Ice Loss on Daily Weather Patterns over North America. J. Clim. 37, 1065–1085 (2024), <u>https://doi.org/10.1175/JCLI-D-23-0389.1</u>
- Ye, K., Woollings, T., Sparrow, S. N., Watson, P. A. G. & Screen, J. A. Response of winter climate and extreme weather to projected Arctic sea-ice loss in very large-ensemble climate model simulations. npj Clim. Atmos. Sci. 7, (2024), <u>https://doi.org/10.1038/s41612-023-00562-5</u>
- Hay, S., Blockley, E., Catto, J. L., Hewitt, H., Lo, Y. T. E., Screen, J. A., Smith, D. M. & Yu, H. The impact of Arctic sea-ice loss on winter weather in the British Isles. Q. J. R. Meteorol. Soc, (2025), accepted
- 36. Smith, D. M., Eade, R., Andrews, M. B., Ayres, H., Clark, A., Chripko, S., Deser, C., Dunstone, N. J., García-Serrano, J., Gastineau, G., Graff, L. S., Hardiman, S. C., He, B., Hermanson, L., Jung, T., Knight, J., Levine, X., Magnusdottir, G., Manzini, E., Matei, D., Mori, M., Msadek, R., Ortega, P., Peings, Y., Scaife, A. A., Screen, J. A., Seabrook, M., Semmler, T., Sigmond, M., Streffing, J., Sun, L. & Walsh, A. Robust but weak winter atmospheric circulation response to future Arctic sea ice loss. Nature Commun. 13, (2022), https://doi.org/10.1038/s41467-022-28283-y
- Screen, J. A., Eade, R., Smith, D. M., Thomson, S. & Yu, H. Net Equatorward Shift of the Jet Streams When the Contribution From Sea-Ice Loss Is Constrained by Observed Eddy Feedback. Geophys. Res. Lett. 49, (2022), <u>https://doi.org/10.1029/2022GL100523</u>
- 38. Hay, S., Catto, J. L., Screen, J. A. & Yu, H. The tug-of-war on the storm tracks between seaice loss and ocean warming is mainly an Atlantic phenomenon. J. Clim. (2025), accepted

- Ye, K., Woollings, T. & Screen, J. A. European Winter Climate Response to Projected Arctic Sea-Ice Loss Strongly Shaped by Change in the North Atlantic Jet. Geophys. Res. Lett., (2023), <u>https://doi.org/10.1029/2022GL102005</u>
- Anderson, Y. A., Perez, J. & Maycock, A. C. Minimal influence of future Arctic sea ice loss on North Atlantic jet-stream morphology. Weather Clim. Dynam. 6, 595–608 (2025), <u>https://doi.org/10.5194/wcd-6-595-2025</u>
- Ronalds, B., Barnes, E. A., Eade, R., Peings, Y. & Sigmond, M. North Pacific zonal wind response to sea ice loss in the Polar Amplification Model Intercomparison Project and its downstream implications. Clim. Dyn. 55, 1779–1792 (2020), https://doi.org/10.1007/s00382-020-05352-w
- Deser, C., Sun, L., Tomas, R. A., & Screen, J. Does ocean coupling matter for the northern extratropical response to projected Arctic sea ice loss? Geophys. Res. Lett. 43, 2149–2157 (2016), <u>https://doi.org/10.1002/2016GL067792</u>
- Hay, S., Priestley, M. D. K., Yu, H., Catto, J. L. & Screen, J. A. The Effect of Arctic Sea-Ice Loss on Extratropical Cyclones. Geophys. Res. Lett. 50, (2023), <u>https://doi.org/10.1029/2023GL102840</u>
- Audette, A. & Kushner, P. J. Role of ocean coupling in the weakening of the extratropical storm tracks from Arctic sea ice loss. Environ. Res. Clim. 4, 025001 (2025), <u>https://doi.org/10.1088/2752-5295/adc828</u>
- Shaw, T. A. & Smith, Z. The Midlatitude Response to Polar Sea Ice Loss: Idealized Slab-Ocean Aquaplanet Experiments with Thermodynamic Sea Ice. J. Clim. 35, 2633–2649 (2022), <u>https://doi.org/10.1175/JCLI-D-21-0508.1</u>
- Riebold, J., Richling, A., Ulbrich, U., Rust, H., Semmler, T. & Handorf, D. On the linkage between future Arctic sea ice retreat, Euro-Atlantic circulation regimes and temperature extremes over Europe. Weather Clim. Dynam. 4, 663–682 (2023), https://doi.org/10.5194/wcd-4-663-2023
- Saurral, R. I., Doblas-Reyes, F. J., Screen, J. A., Catto, J. L., Hay, S. & Yu, H. Western Mediterranean Droughts Fostered by Arctic Sea Ice Loss. J. Clim. 38, 3005–3014 (2025), https://doi.org/10.1175/JCLI-D-25-0066.1
- Hai, L., Zhan, R., Zhao, J. & Wu, B. Spring Barents Sea ice loss enhances tropical cyclone genesis over the eastern North Pacific. Clim. Dyn. 62, 4967–4979 (2024), https://doi.org/10.1007/s00382-024-07145-x
- Ma, W., Chen, G., Peings, Y. & Alviz, N. Atmospheric River Response to Arctic Sea Ice Loss in the Polar Amplification Model Intercomparison Project. Geophys. Res. Lett. 48, (2021), <u>https://doi.org/10.1029/2021GL094883</u>
- 50. Yu, H., Screen, J. A., Hay, S., Catto, J. L. & Xu, M. Winter Precipitation Responses to Projected Arctic Sea Ice Loss and Global Ocean Warming and Their Opposing Influences

over the Northeast Atlantic Region. J. Clim. 36, 4951–4966 (2023), https://doi.org/10.1175/JCLI-D-22-0774.1

- Lewis, N. T., Seviour, W. J. M., Roberts-Straw, H. E. & Screen, J. A. The Response of Surface Temperature Persistence to Arctic Sea-Ice Loss. Geophys. Res. Lett. 51, (2024a), https://doi.org/10.1029/2023GL106863
- Blackport, R. & Fyfe, J. C. Amplified warming of North American cold extremes linked to human-induced changes in temperature variability. Nature Commun. 15, 5864 (2024), <u>https://doi.org/10.1038/s41467-024-49734-8</u>
- Screen, J.A. Arctic amplification decreases temperature variance in northern mid- to highlatitudes. Nature Clim. Change 4, 577–582 (2014), <u>https://doi.org/10.1038/nclimate2268</u>
- Lo, Y. T. E., Mitchell, D. M., Watson, P. A. G. & Screen, J. A. Changes in Winter Temperature Extremes From Future Arctic Sea-Ice Loss and Ocean Warming. Geophys. Res. Lett. 50, (2023), <u>https://doi.org/10.1029/2022GL102542</u>
- Kang, J. M., Shaw, T. A. & Sun, L. Arctic Sea Ice Loss Weakens Northern Hemisphere Summertime Storminess but Not Until the Late 21st Century. Geophys. Res. Lett. 50, (2023), <u>https://doi.org/10.1029/2022GL102301</u>
- 56. Kay, G., Dunstone, N., Smith, D., Dunbar, T., Eade, R. & Scaife, A. Current likelihood and dynamics of hot summers in the UK. Environ. Res. Lett. 15, 094099 (2020), <u>https://doi.org/10.1088/1748-9326/abab32</u>
- Sellevold, R., Lenaerts, J. T. M. & Vizcaino, M. Influence of Arctic sea-ice loss on the Greenland ice sheet climate. Clim. Dyn. 58, 179–193 (2021), <u>https://doi.org/10.1007/s00382-021-05897-4</u>
- Zhang, X., He, B., Bao, Q., Liu, Y., Wu, G., Duan, A., Hu, W., Sheng, C. & Rao, J. The role of Arctic sea ice loss in the interdecadal trends of the East Asian summer monsoon in a warming climate. npj Clim. Atmos. Sci. 7, (2024), <u>https://doi.org/10.1038/s41612-024-00717-y</u>
- Deser, C., Tomas, R. A., Alexander, M. A., & Lawrence, D. M. The seasonal atmospheric response to projected Arctic sea ice loss in the late twenty-first century. J. Clim. 23, 333– 351 (2010), <u>https://doi.org/10.1175/2009JCLI3053.1</u>
- Ye, K., Woollings, T. & Sparrow, S. N. Dynamic and Thermodynamic Control of the Response of Winter Climate and Extreme Weather to Projected Arctic Sea-Ice Loss. Geophys. Res. Lett. 51, (2024b), <u>https://doi.org/10.1029/2024GL109271</u>
- Streffing, J., Semmler, T., Zampieri, L. & Jung, T. Response of Northern Hemisphere weather and climate to Arctic sea ice decline: Resolution independence in Polar Amplification Model Intercomparison Project (PAMIP) simulations. J. Clim. 1–39 (2021), <u>https://doi.org/10.1175/JCLI-D-19-1005.1</u>

- Sigmond, M. & Sun, L. The role of the basic state in the climate response to future Arctic sea ice loss. Environ. Res.: Clim. 3, 031002 (2024) <u>https://doi.org/10.1088/2752-5295/ad44ca</u>
- Dai, A. & Song, M. Little influence of Arctic amplification on mid-latitude climate. Nature Clim. Change 10, 231–237 (2020), <u>https://doi.org/10.1038/s41558-020-0694-3</u>
- Scaife, A. A. & Smith, D. A signal-to-noise paradox in climate science. npj Clim. Atmos. Sci. 1, (2018), <u>https://doi.org/10.1038/s41612-018-0038-4</u>
- Scaife, A. A., Camp, J., Comer, R., Davis, P., Dunstone, N., Gordon, M., MacLachlan, C., Martin, N., Nie, Y., Ren, H., Roberts, M., Robinson, W., Smith, D. & Vidale, P. L. Does increased atmospheric resolution improve seasonal climate predictions? Atmos. Sci. Lett. 20, (2019), <u>https://doi.org/10.1002/asl.922</u>
- 66. Smith, D. M., Dunstone, N. J., Scaife, A. A., Fiedler, E. K., Copsey, D. & Hardiman, S. C. Atmospheric Response to Arctic and Antarctic Sea Ice: The Importance of Ocean– Atmosphere Coupling and the Background State. J. Clim. 30, 4547–4565 (2017), https://doi.org/10.1175/JCLI-D-16-0564.1
- Mudhar, R., Seviour, W. J. M., Screen, J. A., Geen, R., Lewis, N. T. & Thomson, S. I. Exploring Mechanisms for Model-Dependency of the Stratospheric Response to Arctic Warming. J. Geophys. Res. Atmos. 129, (2024), <u>https://doi.org/10.1029/2023JD040416</u>
- Xu, M., Tian, W., Zhang, J., Screen, J. A., Zhang, C. & Wang, Z. Important role of stratosphere-troposphere coupling in the Arctic mid-to-upper tropospheric warming in response to sea-ice loss. npj Clim. Atmos. Sci. 6, (2023), <u>https://doi.org/10.1038/s41612-023-00333-2</u>
- Labe, Z., Peings, Y. & Magnusdottir, G. Warm Arctic, Cold Siberia Pattern: Role of Full Arctic Amplification Versus Sea Ice Loss Alone. Geophys. Res. Lett. 47, (2020), <u>https://doi.org/10.1029/2020GL088583</u>
- Labe, Z., Peings, Y. & Magnusdottir, G. The Effect of QBO Phase on the Atmospheric Response to Projected Arctic Sea Ice Loss in Early Winter. Geophys. Res. Lett. 46, 7663– 7671 (2019), <u>https://doi.org/10.1029/2019GL083095</u>
- Xu, M., Screen, J. A., Tian, W., Zhang, J., Zhang, C. & Yu, H. Influence of Regional Sea Ice Loss on the Arctic Stratospheric Polar Vortex. J. Geophys. Res. Atmos. 129, (2024), <u>https://doi.org/10.1029/2023JD040571</u>
- Walsh, A., Screen, J. A., Sciafe, A. A., Smith, D. M. & Eade, R. Interdependent extratropical atmospheric responses to Arctic sea-ice loss, QBO and ENSO. J. Clim. (2025), submitted
- Screen, J. A. & Francis, J. A. Contribution of sea-ice loss to Arctic amplification is regulated by Pacific Ocean decadal variability. Nature Clim. Change 6, 856–860 (2016), <u>https://doi.org/10.1038/nclimate3011</u>

- 74. Osborne, J. M., Screen, J. A. & Collins, M. Ocean–Atmosphere State Dependence of the Atmospheric Response to Arctic Sea Ice Loss. J. Clim. 30, 1537–1552 (2017), <u>https://doi.org/10.1175/JCLI-D-16-0531.1</u>
- Liang, Y.-C., Kwon, Y.-O., Frankignoul, C., Gastineau, G., Smith, K. L., Polvani, L. M., Sun, L., Peings, Y., Deser, C., Zhang, R. & Screen, J. The Weakening of the Stratospheric Polar Vortex and the Subsequent Surface Impacts as Consequences to Arctic Sea Ice Loss. J. Clim. 37, 309–333 (2024), <u>https://doi.org/10.1175/JCLI-D-23-0128.1</u>
- 76. Simon, A., Gastineau, G., Frankignoul, C., Lapin, V. & Ortega, P. Pacific Decadal Oscillation modulates the Arctic sea-ice loss influence on the midlatitude atmospheric circulation in winter. Weather Clim. Dynam. 3, 845–861 (2022), https://doi.org/10.5194/wcd-3-845-2022
- 77. Cvijanovic, I., Simon, A., Levine, X., White, R., Ortega, P., Donat, M., Lucas, D. D., Chiang, J. C. H., Seidenglanz, A., Bojovic, D., Amaral, A. R., Lapin, V., Doblas-Reyes, F. & Petrova, D. Arctic sea-ice loss drives a strong regional atmospheric response over the North Pacific and North Atlantic on decadal scales. Commun. Earth Environ. 6 (2025), https://doi.org/10.1038/s43247-025-02059-w
- Rosenblum, E., & Eisenman. Sea ice trends in climate models only accurate in runs with biased global warming. J. Clim. 30, 6265-6278 (2017), <u>https://doi.org/10.1175/JCLI-D-16-0455.1</u>
- Armour, K. C., Marshall, J., Scott, J. R., Donohoe, A. & Newsom, E. R. Southern Ocean warming delayed by circumpolar upwelling and equatorward transport. Nature Geosci. 9, 549–554 (2016), <u>https://doi.org/10.1038/ngeo2731</u>
- Lee, Y.-C., Liu, W., Fedorov, A. V., Feldl, N. & Taylor, P. C. Impacts of Atlantic meridional overturning circulation weakening on Arctic amplification. Proc. Natl. Acad. Sci. U.S.A. 121, (2024), <u>https://doi.org/10.1073/pnas.2402322121</u>
- Pithan, F. & Jung, T. Arctic Amplification of Precipitation Changes—The Energy Hypothesis. Geophys. Res. Lett. 48, (2021), <u>https://doi.org/10.1029/2021GL094977</u>
- Hobbs, W., Spence, P., Meyer, A., Schroeter, S., Fraser, A. D., Reid, P., Tian, T. R., Wang, Z., Liniger, G., Doddridge, E. W. & Boyd, P. W. Observational Evidence for a Regime Shift in Summer Antarctic Sea Ice. J. Clim. 37, 2263–2275 (2024), <u>https://doi.org/10.1175/JCLI-D-23-0479.1</u>
- Purich, A. & Doddridge, E. W. Record low Antarctic sea ice coverage indicates a new sea ice state. Commun. Earth Environ. 4, (2023), <u>https://doi.org/10.1038/s43247-023-00961-9</u>
- Raphael, M. N., Maierhofer, T. J., Fogt, R. L., Hobbs, W. R. & Handcock, M. S. A twentyfirst century structural change in Antarctica's sea ice system. Commun. Earth Environ. 6, (2025), <u>https://doi.org/10.1038/s43247-025-02107-5</u>

- Ayres, H. C., Screen, J. A., Blockley, E. W., & Bracegirdle, T. J. The coupled atmosphereocean response to Antarctic sea-ice loss. J. Clim. 35, 4665–4685 (2022), https://doi.org/10.1175/JCLI-D-21-0918.1
- Josey, S.A., Meijers, A.J.S., Blaker, A.T., Blaker, A.T., Grist, J.P., Mecking, J, Ayres, H.C.. Record-low Antarctic sea ice in 2023 increased ocean heat loss and storms. Nature 636, 635–639 (2024).
- Deser, C., Tomas, R. A., & Sun, L. The role of ocean–atmosphere coupling in the zonalmean atmospheric response to Arctic sea ice loss. J. Clim. 28, 2168–2186 (2015), <u>https://doi.org/10.1175/JCLI-D-14-00325.1</u>
- Deng, J. & Dai, A. Sea ice–air interactions amplify multidecadal variability in the North Atlantic and Arctic region. Nature Commun. 13, 2100 (2022), <u>https://doi.org/10.1038/s41467-022-29810-7</u>
- Yu, Q., Wu, B. & Zhang, W. The atmospheric connection between the Arctic and Eurasia is underestimated in simulations with prescribed sea ice. Commun. Earth Environ. 5, (2024), <u>https://doi.org/10.1038/s43247-024-01605-2</u>
- 90. Xu, M., Kang, S., Screen, J. A., Audette, A., Tian, W., Zhang, J., Yu, H. & Zhang, H.: Ocean-atmosphere coupling enhances Eurasian cooling in response to historical Barents-Kara sea-ice loss. npj Clim. Atmos. Sci. (2025), submitted
- England, M. R., Eisenman, I. & Wagner, T. J. W. Spurious Climate Impacts in Coupled Sea Ice Loss Simulations. J. Clim. 35, 7401–7411 (2022), <u>https://doi.org/10.1175/JCLI-D-21-0647.1</u>
- England, M. R., Feldl, N. & Eisenman, I. Sea ice perturbations in aquaplanet simulations: isolating the physical climate responses from model interventions. Environ. Res.: Clim. 3, 045031 (2024), <u>https://doi.org/10.1088/2752-5295/ad9b45</u>
- Lewis, N. T., England, M. R., Screen, J. A., Geen, R., Mudhar, R., Seviour, W. J. M. & Thomson, S. I. Assessing the Spurious Impacts of Ice-Constraining Methods on the Climate Response to Sea Ice Loss Using an Idealized Aquaplanet GCM. J. Clim. 37, 6729–6750 (2024b), <u>https://doi.org/10.1175/JCLI-D-24-0153.1</u>
- 94. Fraser-Leach, L., Kushner, P. & Audette, A. Correcting for artificial heat in coupled sea ice perturbation experiments. Environ. Res.: Clim. 3, 015003 (2023), <u>https://doi.org/10.1088/2752-5295/ad1334</u>
- Semmler, T., Pithan, F. & Jung, T. Quantifying two-way influences between the Arctic and mid-latitudes through regionally increased CO2 concentrations in coupled climate simulations. Clim. Dyn. 54, 3307–3321 (2020), <u>https://doi.org/10.1007/s00382-020-05171-z</u>
- Stuecker, M. F., Bitz, C. M., Armour, K. C., Proistosescu, C., Kang, S. M., Xie, S.-P., Kim, D., McGregor, S., Zhang, W., Zhao, S., Cai, W., Dong, Y. & Jin, F.-F. Polar amplification

dominated by local forcing and feedbacks. Nature Clim. Change 8, 1076–1081 (2018), https://doi.org/10.1038/s41558-018-0339-y

- Simon, A., Gastineau, G., Frankignoul, C., Rousset, C. & Codron, F. Transient Climate Response to Arctic Sea Ice Loss with Two Ice-Constraining Methods. J. Clim. 34, 3295– 3310 (2021), <u>https://doi.org/10.1175/JCLI-D-20-0288.1</u>
- Dai, A., Luo, D., Song, M., & Liu, J. Arctic amplification is caused by sea-ice loss under increasing CO₂. Nat. Commun. 10, 121 (2019), <u>https://doi.org/10.1038/s41467-018-07954-9</u>
- 99. Wijngaard, R. R., van de Berg, W. J., van Dalum, C. T., Herrington, A. R. & Levine, X. J. Arctic temperature and precipitation extremes in present-day and future storyline-based variable resolution Community Earth System Model simulations. (2025), <u>https://doi.org/10.5194/egusphere-2025-1070</u>
- 100. Wills, R. C. J., Herrington, A. R., Simpson, I. R. & Battisti, D. S. Resolving Weather Fronts Increases the Large-Scale Circulation Response to Gulf Stream SST Anomalies in Variable-Resolution CESM2 Simulations. J. Adv. Model Earth Syst. 16, (2024), https://doi.org/10.1029/2023MS004123
- 101. Geen, R., Thomson, S. I., Screen, J. A., Blackport, R., Lewis, N. T., Mudhar, R., Seviour, W. J. M. & Vallis, G. K. An Explanation for the Metric Dependence of the Midlatitude Jet-Waviness Change in Response to Polar Warming. Geophys. Res. Lett. 50, (2023), https://doi.org/10.1029/2023GL105132
- 102. Kochkov, D., Yuval, J., Langmore, I., Norgaard, P., Smith, J., Mooers, G., Klöwer, M., Lottes, J., Rasp, S., Düben, P., Hatfield, S., Battaglia, P., Sanchez-Gonzalez, A., Willson, M., Brenner, M. P. & Hoyer, S. Neural general circulation models for weather and climate. Nature 632, 1060–1066 (2024), <u>https://doi.org/10.1038/s41586-024-07744-y</u>
- 103. Yu, H., Screen, J. A., Xu, M., Hay, S., Qiu, W. & Catto, J. L. Incomplete Arctic Sea-Ice Recovery Under CO2 Removal and Its Effects on the Winter Atmospheric Circulation. Geophys. Res. Lett. 52, (2025), https://doi.org/10.1029/2024GL113541
- 104. Chatterjee, S., Semmler, T., Screen, J., Furevik, T., Selivanova, J., Raj, R. P., Murukesh, N. & Ravichandran, M. Atmosphere–Ocean–Sea Ice Feedbacks Sustain Recent Barents Sea Ice Loss despite Cooler Atlantic Water Inflow. J. Clim. 37, 6519–6532 (2024), https://doi.org/10.1175/JCLI-D-24-0020.1
- 105. Ding, X., Chen, G., Sun, L. & Zhang, P. Distinct North American Cooling Signatures Following the Zonally Symmetric and Asymmetric Modes of Winter Stratospheric Variability. Geophys. Res. Lett. 49, (2022), <u>https://doi.org/10.1029/2021GL096076</u>
- 106. Ball, E., Eunice Lo, Y. T., AG Watson, P., Lavigne, E., Screen, J., Ye, K. & Mitchell, D. No detectable decrease in extreme cold-related mortality in Canada from Arctic sea ice loss. Environ. Res. Lett. 20, 044042 (2025), <u>https://doi.org/10.1088/1748-9326/adc1e6</u>

- 107. Dong, W., Jia, X., Li, X. & Wu, R. Synergistic effects of Arctic amplification and Tibetan Plateau amplification on the Yangtze River Basin heatwaves. npj Clim. Atmos. Sci. 7, (2024), <u>https://doi.org/10.1038/s41612-024-00703-4</u>
- 108. Luo, B., Luo, D., Dai, A., Xiao, C., Simmonds, I., Hanna, E., Overland, J., Shi, J., Chen, X., Yao, Y., Duan, W., Liu, Y., Zhang, Q., Xu, X., Diao, Y., Jiang, Z. & Gong, T. Rapid summer Russian Arctic sea-ice loss enhances the risk of recent Eastern Siberian wildfires. Nature Commun. 15, (2024), <u>https://doi.org/10.1038/s41467-024-49677-0</u>

Figures



Figure 1: Differences in the specified sea ice and SST between selected PAMIP experiments. Differences in the prescribed annual-mean sea ice concentration (SIC) and sea surface temperature (SST) between PAMIP experiments to quantify the effects of (a) past and (b) future Arctic sea-ice loss; (c) past and (d) future Antarctic sea-ice loss; and (e) past and (f) future SST change. The number of models providing each experiment combination and the total number of members, for each year-long time-slice experiment, across all models (publicly available on the Earth System Grid Federation) are provided. The present-day SIC and SST are representative of the period 1979-2008. The pre-industrial and future periods correspond to when the global mean temperature was 0.6°C cooler and 1.4°C warmer than present-day (2°C warmer than preindustrial), respectively.



Figure 2: Contributions to seasonal Arctic warming from various processes and feedbacks. Arctic warming contributions (in K) in the warm half-year (April-September) *vs.* cold half-year (October-March) from the surface albedo (α , red), water vapor (q; blue), Planck (PL'; black), lapse rate (LR; green) and cloud (C; cyan) feedbacks, and changes in oceanic heat release (- Δ OHU; orange) and atmospheric energy convergence (Δ (- ∇ ·F_A); maroon) in response to (a) past and (b) future SST warming, and (c) past and (d) future Arctic sea-ice loss. Adapted from Jenkins et al. (2024)²¹.



Figure 3: Schematic representation of the mechanisms of the jet shift in response to Arctic seaice loss. Features depicted in black represent the climatological state whereas those shown in colour represent the response to sea-ice loss. See Smith et al. (2022)³⁶ for further details.



Figure 4: Regional effects of Arctic sea-ice loss. Visual summary of the predominantly wintertime effects of Arctic sea-ice loss across the Northern Hemisphere, as simulated in the PAMIP experiments. The assessed confidence is based on expert judgement, considering the level of consistency between studies/models/experiments, knowledge of the mechanisms involved, and evidence that the simulated response is representative of that in the real world. The locations of features are approximate. The supporting references are examples and are not intended to provide a comprehensive list of all studies that have reported on each effect.



Figure 5. Distinct responses to sea-ice loss and SST warming. Multimodel-mean winter near-surface air temperature response to (a) future Arctic sea-ice loss and (b) future SST warming, and (c) their relative signs and magnitudes. Hatching denotes statistical significance at the 90% confidence level. In (c), yellow and orange colors denote the responses reinforce each other, whereas blue and red colors denote the responses oppose each other, and stippling shows where the response to sea-ice loss is of greater magnitude than that to SST warming. (d-f) As a-c, but for precipitation. (g-i) As a-c, but for 850 hPa westerly wind. Adapted from Yu et al. (2024)²⁸.



Figure 6. Storylines of projected climate change. (a-c) Summed multimodel winter near-surface air temperature responses to future Arctic sea-ice loss and SST warming, for three different storylines: low sea-ice sensitivity (1.5 million square kilometers of sea-ice loss per degree Celsius of global-mean SST warming), medium sea-ice sensitivity (2.7 million km²/°C) and high sea-ice sensitivity (3.8 million km²/°C). The low and high sensitivity storylines are calculated by linearly scaling the PAMIP responses prior to their summation; the medium sea-ice sensitivity storyline uses the unscaled PAMIP responses. (d) The difference between the high and low sensitivity storylines. (e-h) As a-d, but for precipitation. (i-l) As a-d, but for 850 hPa westerly wind. Hatching denotes statistical significance at the 90% confidence level. Numbers in the lower left corner denote Arctic amplification in a-c, Arctic-mean precipitation change in e-g, and the North Atlantic jet latitude change in i-k. Adapted from Yu et al. (2024)²⁸ and Hay et al. (2025)³⁵.