

Projected Changes in Both Mean Climate and Climate Variability Drive Substantial Increases in Extreme Fire Weather in the Western United States

DANIELLE TOUMA^{a,b} AND CLARA DESER^b

^a *Jackson School of Geosciences, Institute for Geophysics, The University of Texas at Austin, Austin, Texas*

^b *Climate and Global Dynamics, NSF National Center for Atmospheric Research, Boulder, Colorado*

(Manuscript received 5 February 2025, in final form 30 April 2026, accepted 20 May 2026)

ABSTRACT: Wildfire frequency, extent, and duration in the western United States are projected to increase throughout the twenty-first century as dry and warm conditions become more frequent, widespread, and persistent. However, there is limited knowledge about how changes in mean climate and changes in climate variability, individually and together, contribute to these increases. To disentangle these effects, we use a 100-member ensemble of climate simulations produced with the Community Earth System Model, version 2, under historic and shared socioeconomic pathway (SSP) 3-7.0 forcing from 1980 to 2100. Using the Canadian forest fire weather index (FWI) to quantify fire-related meteorological conditions in the simulations, we select extreme FWI thresholds relative to a baseline distribution centered at 1980 (“fixed threshold”) and an evolving distribution (“moving threshold”) to identify spatiotemporally connected extreme fire weather events. By 2100, the frequency, area, and number of event days increase when considering a fixed threshold, and events are up to 4°C warmer on average. Moreover, events over the Pacific Coast expand northwestward, while those over the Four Corners region expand westward and northeastward. Changes in event frequency are small and not significant when considering a moving baseline but become spatiotemporally more connected and widespread, reflecting nonthermodynamically driven changes. Increases in the mean of maximum temperature, as well as changes in both the variability and mean of relative humidity, drive the largest increases in extreme fire weather event area and number of event days. By quantifying how changes in the mean and variability of climate variables impact wildfire conditions in the western United States, our study has implications for increasing resiliency to wildfire risk under climate change.

KEYWORDS: Climate change; Climate variability; Ensembles; Forest fires

1. Introduction

Wildfires in recent years have become more synchronous, occurring in different parts of the North American continent simultaneously and compounding the impacts felt in the region (Abatzoglou et al. 2021). For example, in 2023, unprecedented wildfires in Canada were burning from east to west across the country simultaneously, straining the capacity for coordinated firefighting and leading to poor air quality in much of the United States and Canada (Jones et al. 2024; Kirchmeier-Young et al. 2024). Similarly, in 2020, the western United States experienced widespread wildfire activity—while the number of fires was in the normal historic range, the burn area and the number of very large fires were the highest recorded, leading to 40 fatalities and \$20 billion (U.S. dollars) in losses. Moreover, in much of the western United States, wildfire area and severity have increased over the last few decades (Jones et al. 2022). Specifically, increases in the total burned area and the number of large wildfires have been attributed to anthropogenically

forced anomalous warming and drying in recent years (Jain et al. 2022; Juang et al. 2022; Abatzoglou et al. 2025) and over a century of forest fire suppression (Andela et al. 2017; Jones et al. 2022; Hagmann et al. 2021). The damages felt by society due to these increases in wildfire area have also risen, with larger populations and more resources being exposed to wildfire hazards (Burke et al. 2021). Moreover, these larger and more frequent wildfires have exacerbated populations’ exposure to smoke and postfire hydrologic hazards, such as debris flows and channel sedimentation (Oakley 2021). These impacts have both short-term and long-term impacts on hydrology (Collar et al. 2022; Williams et al. 2022), livelihoods (Lawrence et al. 2022), and public health (Burke et al. 2021).

Many studies have assessed trends in fire weather conditions—dry, warm, and windy conditions—using fire weather indices such as the Canadian forest fire weather index (FWI) and the McArthur forest fire danger index (FFDI) and their impact on the likelihood, frequency, and area of wildfires in recent years. Observed increases in extreme fire weather events, usually defined using local, extreme (95th or 99th) percentiles of the historical distribution of a fire weather index, lead to increases in burned forest area and higher frequency of large wildfires globally and in the western United States (Abatzoglou and Williams 2016; Goss et al. 2020; Jain et al. 2022; Jones et al. 2022). These increases largely stem from increases in maximum temperature and decreases in relative humidity prior to and during wildfire seasons and have been linked to an increase in the frequency and intensity of atmospheric ridging in the northeast Pacific Ocean (Sharma et al. 2022). The southwest United States has also been impacted by “failed” North American

 Denotes content that is immediately available upon publication as open access.

 Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/JCLI-D-25-0077.s1>.

Corresponding author: Danielle Touma, danielle.touma@utexas.edu

DOI: 10.1175/JCLI-D-25-0077.1

© 2026 American Meteorological Society. This published article is licensed under the terms of the default AMS reuse license. For information regarding reuse of this content and general copyright information, consult the AMS Copyright Policy (www.ametsoc.org/PUBSReuseLicenses).

monsoon seasons as well as delays in monsoon onsets, leading to longer-than-usual dry conditions and subsequently higher wildfire activity (Cook and Seager 2013; Hoell et al. 2022). Changes in these conditions across the western United States are largely driven by increases in global greenhouse gas emissions and localized reductions in aerosol emissions (Abatzoglou and Williams 2016; Kirchmeier-Young et al. 2017; Touma et al. 2021). In future years, projected increases in greenhouse gas emissions and continued reductions in aerosol emissions are expected to lead to further increases in frequencies and durations of extreme fire weather conditions, leading to higher risks of wildfires and causing the frequency of extreme fire weather conditions to emerge well beyond the expected historic frequency, with fire weather conditions reaching a “new normal” by the end of the twenty-first century (Abatzoglou et al. 2019; Touma et al. 2021, 2022).

Several studies have used large-ensemble Earth system model (ESM) simulations to quantify the trends in fire weather conditions in future climate projections. There is a large consensus that fire weather conditions are projected to increase over much of the globe when accounting for both model discrepancies (Abatzoglou et al. 2019; Bui et al. 2024) and irreducible internal variability (Kirchmeier-Young et al. 2017; Touma et al. 2021). Given systemic biases in ESMs, especially in the representation of the FWI (Gallo et al. 2023; Touma et al. 2021) and the driving variables (Fasullo 2020; Simpson et al. 2024), the general approach is to use upper tails of the FWI distribution to define “extreme” fire weather within each ESM.

Here, we compare the more traditional choice of using “fixed” historic or preindustrial distributions of fire weather indices to calculate upper-tail thresholds to describe extreme fire weather conditions to the choice of using a moving-window threshold (hereafter “moving threshold”), which captures the time-evolving distribution of fire weather conditions. In recent marine heat wave literature, it has been argued that using a fixed baseline can obscure local and short-term drivers of these extremes in future scenarios due to the saturation of marine heat wave occurrences caused by a warming climate (Smith et al. 2025; Capotondi et al. 2024; Deser et al. 2024). For terrestrial heat waves, studies have compared changes in heat waves defined using a moving baseline to those using a fixed baseline to separate thermodynamic from nonthermodynamic effects on changes in heat wave area and duration (Skinner et al. 2025; Vogel et al. 2020). Using a moving threshold can also reveal additional criteria for adaptation measures needed to overcome changes in extreme climate events, beyond those driven by mean changes alone (Amaya et al. 2023). In our study, a moving threshold would ensure that the frequency of fire weather days for a specific location remain constant through time; however, the intensity and area of events could still change through changes in the variability beyond the extreme threshold. Moreover, spatial or temporal clustering or connectivity of extreme fire weather events could also increase, due to nonthermodynamic changes. This could lead to greater spatially compounding risks with multiple wildfires occurring simultaneously or in succession, resulting in increased burned area, greater smoke exposure from wildfire emissions, and the preconditioning of larger areas to postfire hydrologic hazards (Raymond et al. 2020; Zscheischler et al. 2020).

Few studies have assessed how changes in the variability of the climate system, and not just the mean of the climate, has impacted wildfire risk under anthropogenic forcing. By doing so, changes in the variability are assumed to be negligible or part of the “noise” of our climate system—however, this could lead to the underestimation of the impact of anthropogenic forcing on wildfire risk. Zhuang et al. (2021) found that changes in atmospheric circulation, or natural variability, explained one-third of the observed vapor pressure deficit increase over the western United States in recent decades. While their study leveraged multimodel simulations from the CMIP6 archive, single-model initial-condition large ensembles could provide a more robust estimate of these influences by overcoming uncertainties from model physics (Deser et al. 2020; Zhuang et al. 2021).

In this study, we use the 100-member Community Earth System Model, version 2, Large Ensemble (CESM2-LE) to assess projected changes under shared socioeconomic pathway (SSP) 3-7.0 (Rodgers et al. 2021) in the Four Corners [Utah (UT), Colorado (CO), New Mexico (NM), and Arizona (AZ)] and the Pacific Coast [California, Oregon (OR), and Washington] regions, which have been experiencing increasing trends in large and severe wildfires (Abatzoglou et al. 2025). We show how extreme fire weather events respond to anthropogenic climate change in terms of their frequency, number of event days, area, and intensity during the height of the western U.S. fire season, June–October (JJASO). We address two main research gaps in understanding wildfire risk in the western United States. First, we use both a fixed- and moving-window period to calculate location-specific extreme percentile thresholds and identify spatiotemporally connected grid points as extreme fire weather events. By using these two types of thresholds, we disentangle the thermodynamic drivers of changes in extreme fire weather events from other nonthermodynamic drivers and provide insight on fire-related adaptation needs in these regions. Second, through the development of a novel framework, we isolate the effects of changes in the mean and variability of climate drivers on changes in the number of event days, area, frequency, and intensity of these events. This allows new insight into how changes in climate variability, in addition to changes in the mean climate, will influence wildfire risk.

2. Data and methods

a. CESM2-LE simulations

We use the CESM2-LE simulations to quantify historic and future fire weather. As demonstrated in previous studies (e.g., Kirchmeier-Young et al. 2017; Touma et al. 2021, 2022), using a large ensemble allows robust assessment of changes in extreme fire weather events under anthropogenic forcing as well as internal variability. The CESM2-LE consists of 100 members and has daily resolution data publicly available, making it a unique and unprecedented dataset to understand future projections of extreme fire weather events.

The CESM2 model is a fully coupled atmosphere, ocean, land, and sea ice model on a nominal 1° grid. The 100 members of the CESM2-LE are simulated under historic (1850–2014)

and SSP3-7.0 (2015–2100) scenarios (Rodgers et al. 2021). The SSP3-7.0 scenario has both high greenhouse gases and aerosol emissions—this was chosen for the CESM2-LE to enable the detection and quantification of forced changes in the mean and natural variability (Rodgers et al. 2021), making it suitable for our study. The ensemble members are initialized through a combination of macro- and microperturbations, where microperturbations are introduced using different Atlantic meridional overturning circulation (AMOC) states, and microperturbations are introduced using a random perturbation to the atmospheric temperature field (Rodgers et al. 2021). The memory of the initial conditions is lost within a few decades (Deser et al. 2025). We note that all 100 members use identical forcing, except for the biomass burning aerosols. Half of the members follow the Coupled Model Intercomparison Project phase 6 (CMIP6) protocol for biomass burning, which consists of low-pass-filtered (11-yr smoothing) time series except over the period 1990–2020 when high-frequency satellite-based measurements are available (Van Marle et al. 2017). This introduces a discontinuity in temporal variance of biomass burning aerosols, which in turn has been shown to produce a rectified effect on climate conditions over high northern latitudes in association with Arctic sea ice feedbacks (e.g., DeRepentigny et al. 2022). For this reason, the other half of the members use 11-yr smoothed biomass burning time series over the entirety of the simulations [see Rodgers et al. (2021) for additional information]. We find no discernable differences between the two halves of the CESM2-LE members for our analysis and, therefore, consider all 100 members together.

b. Canadian forest FWI

The Canadian forest FWI system (CFFWIS) consists of three moisture codes and three fire behavior indices and was empirically derived by Van Wagner (1987). The FWI itself, which is part of the CFFWIS, is a numeric rating of fire intensity and accounts for the moisture codes and other indices in the system, and its calculation is described in Dowdy et al. (2009). FWI is widely used, both operationally and in understanding the impact of climate change on wildfire risk (Abatzoglou et al. 2019; Touma et al. 2021, 2022, 2023; Jain et al. 2022). We use a modified version of the FWI to quantify fire weather conditions in the simulations to overcome the lack of subdaily data availability. The FWI is calculated for the CESM2-LE using surface daily precipitation (CESM2 variable name: PRECT, abbreviation: pr), maximum temperature (TREFHTMX, tasmax), relative humidity (RHREFHT, hurs), and surface wind speed (U10, sfcWind). We first calculate the three moisture codes that reflect conditions for three depths of fuel and time scales. Each moisture code has a fuel drying and wetting (or rainfall) phase that is calculated sequentially. The fine fuel moisture code (FFMC) reflects moisture levels for shaded litter fuels and on daily time scales and is calculated using tasmax, hurs, and sfcWind for fuel drying and pr for fuel wetting. The Duff moisture code (DMC) reflects moisture levels for decomposed organic material and on monthly time scales and is calculated using pr for fuel wetting and tasmax and hurs for fuel drying. The drought code (DC) reflects moisture levels for deep litter

and soil and is computed using pr for the rainfall phase and tasmax for the drying phase. Using the FFMC and sfcWind, the initial spread index (ISI) is then calculated to represent the fire spread rate, and using the DMC and DC, the build-up index (BUI) is calculated to represent the potential heat release and severity of fire. Last, the ISI and BUI are used to calculate the FWI and represents overall fire danger. All moisture codes and indices are unitless.

We compare the FWI calculated from the CESM2-LE for JJASO to those calculated from observationally based grid-MET variables (Abatzoglou 2013) over the western United States from 1983 to 2024. We show that the mean FWI is generally biased low in CESM2-LE compared to the mean grid-MET FWI and that the 99th percentile of the FWI is biased high (Fig. 1 in the online supplemental material). While the CESM2-LE captures some large-scale spatial variations in FWI mean and extreme values (e.g., over the Rocky Mountains), smaller-scale variations are not reflected, such as over the Sierra Nevada or the Cascade Mountain Ranges. The bias in the mean FWI is similar to that found in Gallo et al. (2023), who calculated FWI for 16 CMIP6 models (excluding CESM2) and found that the multimodel mean of FWI was biased low for the western United States. We note that the biases are largely systematic over the region, giving us more confidence in using model-based thresholds and analyzing time periods in model data only (i.e., we do not compare future projections with observations).

c. Extreme event definition and baselines

To identify extreme fire weather events, we use the location-specific 99th percentile of the FWI across the full 100-member ensemble. Previous studies have shown that the 95th percentile and above of FWI is well correlated with large and severe wildfires in the western United States (Goss et al. 2020; Jain et al. 2024). While we do not employ a sensitivity analysis to explore the robustness of our results to the 99th percentile threshold, previous studies have shown that using a lower threshold (e.g., 95th percentile) results in similar relative spatial patterns of extreme fire weather events, with more frequent events but smaller relative increases in future periods (Touma et al. 2022) and earlier emergence of future extreme fire weather frequency above the historic variability (Abatzoglou et al. 2019). By using the 99th percentile, our aim is to find events that would have severe repercussions on a region. We use the full ensemble to calculate the 99th percentile to robustly account for internal climate variability when estimating extreme values. We use two baseline types to calculate the 99th percentile threshold and associated FWI variable (pr, tasmax, hurs, and sfcWind) anomalies: 1) a fixed baseline 35-yr period centered around 1980 and 2) a “moving” baseline 35-yr period centered around each year in the time series. We use all calendar day values of the FWI to calculate the percentile threshold value. Using a 35-yr period allows us to capture interannual-to-decadal variability of the FWI and reduce the sensitivity to the exact set of years used to define the baseline. The threshold using the fixed baseline is constant in time, while that with the moving baseline changes from year to




FIG. 1. (a) Example FWI time series for one grid point near Los Angeles and one ensemble member (1001.001), with fixed and moving 99th percentile thresholds and identified exceedances. (b) Annual number of exceedances calculated from (a) fixed and moving thresholds. (c) Example of a 6-day extreme fire weather event over the Pacific Coast region. The shading represents the value of FWI, and the black dots represent the identified spatiotemporally connected event. The event starts when at least one grid point in the regional boundary of the Pacific Coast region (shown in Fig. 3) exceeds the local threshold, and the event ends when there are no longer any grid points that are spatiotemporally connected to that event within the region boundary.

year. By using a moving baseline, we can begin to isolate the long-term, thermodynamically driven trends in the characteristics of extreme fire weather events from the short-term variations of these characteristics. These two approaches bookend the range of potential adaptation to long-term changes—the fixed threshold assumes no adaptation, while the moving threshold assumes continued adaptation to the mean climate. Figures 1a and 1b show an example of these thresholds and identified exceedances for a grid point near Los Angeles.

d. Event identification

For each ensemble member and each threshold type, we identify grid points that meet or exceed the 99th percentile FWI threshold for each time step. We use a framework with image processing functions from the SciPy Python package (Virtanen et al. 2020) to identify spatiotemporally connected grid points that exceed the extreme threshold. We first spatially subset the FWI dataset from 0° to 90°N and from 0° to 180° to increase computational efficiency but ensure that events are not truncated spuriously. We then create a binary dataset, where grid points that meet or exceed the extreme threshold are equal to 1, and those that fall below the threshold are equal to 0. We then identify spatiotemporally connected grid points that are equal to 1—for spatial connectivity, grid points must be connected by their edges or vertices, and for temporal connectivity, at least one grid point within the event must be equal to 1 on two continuous days. Events can be as small as one grid cell and as short as 1 day. We identify events continuously for the full analysis period for all calendar days. The duration and cumulative area of each event is calculated, as well as the corresponding FWI values and FWI input variable anomalies throughout the event.

We identify events for eight U.S. regions (seven individual states, with California separated into northern and southern portions along 37°N) and two regional groupings that have historically experienced large and severe wildfires. These are as follows: Pacific Coast group, comprised of Southern and Northern California, Oregon, and Washington; Four Corners group, comprised of New Mexico, Arizona, Utah, and Colorado. We then find events that have at least one grid point within the bounds of a region or region group of interest for the full duration of the event. If an event in the full event dataset “leaves” the region or region group and then “returns,” it is considered two distinct events. For region groups, we subset the dataset for the full region group simultaneously. This ensures that we do not double count events when events span multiple regions within the group simultaneously. We show an example of a 6-day event that occurred in the Pacific Coast region in Fig. 1c.

e. Event analysis

For each event, we calculate its total duration (number of event days) and area within the region and intensity (event-maximum FWI). We also calculate the anomalies of pr, tasmax, hurs, and scfWind during the event using the 30-day, 35-yr fixed baseline (1963–97) mean for events identified using a fixed threshold and using the 30-day, 35-yr moving baseline mean for those identified using a moving threshold. The 30-day moving mean allows us to account for the seasonal patterns of the FWI input variables. We specify which type of baseline is being used when describing any relevant results.

We create annual distributions of event characteristics including event frequency, number of events days, area, and intensity, as well as for the FWI input variables, for each season

and region by first averaging over all events in each ensemble member in each season and region and then grouping those averages together to create a distribution (100 values representing an ensemble member each). We summarize these time-evolving distributions using the 25th, 50th, and 75th percentiles or the interquartile range (IQR).

We also create composites for each period and each ensemble member for gridpoint- and event-specific values of the number of events, annual number of event days, area, and intensity, as well as pr , $tasmax$, $hurs$, and $sfcWind$ anomalies. We also show vapor pressure and vapor pressure deficit (VPD) anomalies to decouple temperature changes from those in relative humidity. Because vapor pressure and VPD are not available CESM2-LE output variables, we calculate saturation vapor pressure (e_s):

$$e_s = 6.11 \times 10^{(7.5 \times tas / 237.3 + tas)}, \quad (1)$$

where tas is the near-surface air temperature ($^{\circ}\text{C}$) and e_s (hPa). We then calculate actual vapor pressure (e) in hPa:

$$e = e_s \times hurs / 100. \quad (2)$$

We then calculate VPD as the difference between e_s and e .

f. Period definitions

We use three time periods to summarize extreme fire weather event characteristics over time. These are a historic period (1980–2014), an early future period (2015–50), and a late future period (2051–82). We end the last period in 2082 because we use 35-yr moving windows to calculate the moving thresholds and means. We calculate changes in the characteristics of fire weather events for the two future periods relative to the historic period. Additionally, we show time series of extreme fire weather event characteristics for the eight regions, with changes computed from the historic period.

g. Isolating the impact of changes in the mean and variability of FWI variables

We isolate the effect of changes in pr , $tasmax$, $hurs$, and $sfcWind$ on the characteristics of extreme fire weather events. We create new “mapped” (m) time series for each variable and ensemble member (n) (pr_m^n , $tasmax_m^n$, $hurs_m^n$, and $sfcWind_m^n$) using quantile mapping to restrict the whole time series to the 100-member monthly distribution in the fixed reference period using the following steps (see Fig. 2 and supplemental Fig. 2). For each day of the time series, we first find the percentile of the value of the variable within its 100-member, 30-day, 35-yr moving window distribution. We then find the corresponding value of that percentile in the 30-day, 35-yr, 100-member distribution of the fixed baseline period centered around 1980 (Fig. 2). By mapping the FWI variable time series to the fixed period, we maintain the relative day-to-day variability and spatial patterns between the mapped variable and the other unmapped variables but ensure that the variability and mean of the time series are constrained to that of the historic distribution. We then calculate four mapped FWI time series for each ensemble member using one of the newly mapped variables and keeping the other variables at their original values




FIG. 2. Schematic of quantile mapping for maximum temperature for 15 Jun 2040 for a grid point near Los Angeles, CA, for ensemble number 1011.001 using its 100-member future distribution (1–30 Jun, 2023–57) and the 100-member historic (1–30 Jun 1963–97) distribution. The historic period is captured using a 35-yr period centered around 1980, while the future distribution is captured using a 35-yr period centered around 2040. The black line and markers represent the future distribution and values, respectively, and the orange lines and markers represent the historic distribution and values, respectively. Steps 1 and 2: Using the $tasmax_o$ value from the future distribution, find the percentile value; steps 3 and 4: Using that percentile value, find $tasmax_m$ from the historic distribution.

(o) [e.g., $FWI_{pr,m}^n = f(pr_m^n, tasmax_o^n, hurs_o^n, sfcWind_o^n)$]. Using these four mapped FWI time series for each ensemble member, we identify and quantify the characteristics of extreme fire weather events as previously described. These events and their characteristics are considered to represent events in which one of the variables is not impacted by forced changes in the mean and variability. We then compare the event characteristics using the mapped and original FWI time series to assess the contribution of changes in the full distribution (i.e., mean and variability) of each variable to changes in extreme fire weather events.

We also create new time series with only forced changes in the mean removed for each variable (see supplemental Fig. 2). By comparing to the original time series, we isolate the impact of forced changes in the mean of each variable on extreme fire weather event characteristics. For each variable, we calculate forced changes in the mean using the difference between the moving 30-day, 35-yr, 100-member mean and the fixed (1980) 30-day, 35-yr, 100-member mean. We then remove these forced changes in the mean from each respective time series for each ensemble member to create four new “detrended” (d) time series (pr_d^n , $tasmax_d^n$, $hurs_d^n$, and $sfcWind_d^n$). As before, we use these new time series to calculate the FWI four additional times, each time using one of the new time series with the mean removed for one of the variables, while all other variables use their original time series [e.g., $FWI_{pr,d}^n = f(pr_d^n, tasmax_o^n, hurs_o^n, sfcWind_o^n)$]. Using these four new FWI time series for each ensemble member, we identify and quantify the characteristics of extreme fire weather events as previously described. These events and their




FIG. 3. Total number of event days per year over each grid point in the (a),(d),(g),(j) historic period (1980–2014) and change in the number of event days per year over each grid point in (b),(c),(e),(f),(h),(i),(k),(l) future periods using (a)–(f) fixed and (g)–(l) moving threshold for events fully or partially located in the (a)–(c),(g)–(i) Pacific Coast and in the (d)–(f),(j)–(l) Four Corners regions in JJASO. For the historic period, all shaded locations have at least 0.66 event days per year, meaning that at least two-thirds of the ensembles show 1 event day per year on average. Similarly, for the future period changes, all shaded locations have an increase or decrease of at least 0.33 event days per year. Stippling indicates that the difference from the historic period is not statistically significant (p value ≤ 0.05).

characteristics are considered to represent events in which one of the variables is not impacted by forced changes in the mean. The difference between the quantile mapped and the detrended extreme fire weather events is used to estimate the effect of changes in the variability on the seasonal average of that variable (see supplemental Fig. 2).

Last, we isolate the role of changes in vapor pressure mean and variability on changes in extreme fire weather events. We use the vapor pressure e calculated using Eqs. (1) and (2) (methods, section 2e) and create two new time series: vapor pressure mapped to its historic distribution (e_m) and detrended vapor pressure by removing the changes in the mean (e_d). We then recalculate hurs using these two new time series hurs_vp_m^n and hurs_vp_d^n . Then, we recalculate FWI for each ensemble member using these new time series for hurs and identify and quantify the characteristics of extreme fire weather events as previously described. This allows us to understand the role of changes in vapor pressure separate from those in temperature due to their competing influences on relative humidity.

3. Results

a. Historic characteristics of extreme fire weather events

Figure 3 shows composite maps of the number of extreme fire weather event days per year for the Pacific Coast and Four Corners regions using fixed and moving thresholds for JJASO—the height of the wildfire season in the western United States. During the historic period, the fixed baseline shows similar patterns to the moving baseline though with smaller magnitudes because the time period used for the fixed

threshold is slightly earlier than the moving threshold centers used in the composite maps (1963–97 vs 1980–2014) (Figs. 3a,d,g,j). We find that events that are partially or fully located in the Pacific Coast region tend to occur more frequently on the eastern edge of the region (>10 event days per year) relative to coastal areas (~ 8 event days per year) and tend to extend further eastward into Montana, Idaho, Nevada, and Arizona (~ 5 event days per year), southward into Baja California, and northward into western Canada (Figs. 3a,g). Four Corners events largely occur in Arizona and Utah (>12 event days per year), with fewer event days occurring in Colorado and New Mexico, especially across the Rocky Mountains. Similarly, extreme fire weather events spread toward the north, reaching Idaho and Wyoming, and toward the west, reaching into Nevada and California (Figs. 3d,j). We note that because the thresholds are based on 35-yr periods centered on either the fixed threshold year or the moving threshold year and using the full 100-member ensemble, there is some variability in terms of the expected exceedances, with number of event days being higher than the expected 1% of days per year ($3.65 \text{ days yr}^{-1}$).

b. Projected changes in extreme fire weather event characteristics

Extreme fire weather events become more frequent, last longer, and cover larger areas by the end of the twenty-first century when using the fixed threshold compared to the moving threshold (Figs. 3 and 4). In the Pacific Coast, events tend to become more frequent in the eastern parts of the region and more widespread toward the north and east, with up to an additional ~ 15 event days per year reaching the Midwest




FIG. 4. (a)–(d), (m)–(p) Frequency, (e)–(h), (q)–(t) duration, and (i)–(l), (u)–(x) state-bound area of extreme fire weather events using a fixed (coral) and moving (teal) 99th percentile threshold for events in the (a)–(l) Pacific Coast and (m)–(x) Four Corners regions for JJASO.

and an additional ~ 21 event days per year reaching western Canada in the late future period (Fig. 3c). The largest increases occur in Washington, with approximately two more events per year, which also last two more days on average and cover $20\,000\text{ km}^2$ more in the late future compared to the historic period (Figs. 4d,h,l). In the southern part of the region, increases in event duration and area are still significant but are slightly reduced in magnitude, with the smallest increases occurring in Southern California (Figs. 4e,i). Events tend to become more connected and extend in all cardinal directions in the Four Corners region in the future compared to the historic period, with the largest spread toward southern Idaho and Nebraska (>15 more event days per year; Figs. 3e,f). Here, we find that extreme fire weather event duration and

area increase similarly across all four states—events are approximately 2 days longer and cover approximately $10\,000\text{ km}^2$ more than in the historic period (Figs. 4q–x). Interestingly, increases in event area and extent are constrained by the Southern Rocky Mountain Range but not by the Middle or Northern/Canadian Rocky Mountain Ranges for both the Pacific Coast and Four Corners regions (Fig. 3).

While the annual frequency of event days for each location using the moving threshold is explicitly set by the threshold (1% of days), the duration and spatial connectivity of the events are not explicitly bounded. Specifically, this 1% of event days could occur more closely together in both space and time due to changes in the spatial or temporal variability of the FWI and, therefore, could change the area and




FIG. 5. (a),(d),(g),(j),(m),(p),(s),(v) Composite maps of precipitation (mm day^{-1}), maximum temperature (K), relative humidity (%), and wind speed (m s^{-1}) anomalies during extreme fire weather events using (a),(g),(m),(s) 30-day, 35-yr fixed mean for the fixed threshold events and (d),(j),(p),(v) 30-day, 35-yr moving mean for the moving threshold events in the Pacific Coast region in JJASO for the historic period (1980–2014). (b),(c),(e),(f),(h),(i),(k),(l),(n),(o),(q),(r),(t),(u),(w),(x) Composite maps showing the changes in these fire weather variables in the two future periods for the fixed and moving thresholds for 2015–50 and 2050–82. The composites are created by averaging the variable anomalies over the area and duration of all events across all ensemble members in each period. Stippling indicates that the difference from the historic period is not statistically significant ($p \text{ value} \leq 0.05$).




FIG. 6. Time series of (a)–(d) precipitation, (e)–(h) maximum temperature, (i)–(l) relative humidity, and (m)–(p) wind speed anomalies during extreme fire weather events in JJASO in (a),(e),(i),(m) Southern CA, (b),(f),(j),(n) Northern CA, (c),(g),(k),(o) OR, and (d),(h),(l),(p) WA using the fixed mean and threshold (coral) and the moving mean and threshold (teal).

duration of events. We see evidence of small but robust increases in the number of event days, generally occurring to the north and east of the Pacific Coast region and to the northeast and northwest of the Four Corners region (Figs. 3h,i). While these changes seem small and negligible compared to a fixed threshold, they point to changes in areas and durations of events that are not reflected in the thermodynamic shift of the FWI distribution and instead point to the effects of changes in the FWI distribution that may be nonthermodynamic. Specifically, the spread of events beyond the specified region shown when using the moving baseline is likely driven by dynamic processes.

c. Meteorological conditions underlying future changes in extreme fire weather events

In the Pacific Coast region, extreme fire weather events in the historical period are characterized by positive anomalies in maximum temperature (up to 10 K) and wind speed (up to 2 m s^{-1}) and negative anomalies in relative humidity (up to -25%) and precipitation (up to -2 mm day^{-1}) compared to the historical 1963–97 30-day mean (see methods; Figs. 5a,d,g,j,n,p,s,v). In the late future period, extreme fire weather events identified

using a fixed threshold show an increase in maximum temperature anomalies by up to 4 K throughout the Pacific Coast region compared to the historic period (Figs. 5i and 6e–h). Changes in other FWI input variables are more spatially heterogeneous: Precipitation anomalies tend to become less negative over California and more negative over Oregon and Washington; relative humidity anomalies tend to become less negative over the coast ($+4\%$) and more negative over the interior of the region (-4%); and wind speeds tend to be about 0.5 m s^{-1} slower over the northern half of the region but unchanging in the southern part of the region (Figs. 5b,c,n,o,t,u and 6a–d,i–p). The changes in these input variables during extreme fire weather events are more acute than when considering all days in JJASO (supplemental Fig. 3). We also note that, while not included in the calculation of the FWI, actual vapor pressure anomalies increase along with VPD anomalies, reflecting similar changes in relative humidity inland but opposite changes on the coast (supplemental Fig. 4).

For the moving threshold case, we employed a moving 35-yr mean of the FWI input variables from which to calculate the anomalies (see methods) to allow for consistency in the context




FIG. 7. (a),(d),(g),(j),(m),(p),(s),(v) Composite maps of precipitation (mm day^{-1}), maximum temperature (K), relative humidity (%), and wind speed (m s^{-1}) anomalies during extreme fire weather events using (a),(g),(m),(s) 30-day, 35-yr fixed mean for the fixed threshold events and (d),(j),(p),(v) 30-day, 35-yr moving mean for the moving threshold events in the Four Corners region in JJASO for the historic period (1980–2014). (b),(c),(e),(f),(h),(i),(k),(l),(n),(o),(q),(r),(t),(u),(w),(x) Composite maps showing the changes in these fire weather variables in the two future periods for the fixed and moving thresholds for 2015–50 and 2050–82. The composites are created by averaging the variable anomalies over the area and duration of all events across all ensemble members in each period. Stippling indicates that the difference from the historic period is not statistically significant (p value ≤ 0.05).

of the threshold that is used to define the events. For these events, future increases in temperature anomalies during extreme fire weather events are limited to <1 K (Figs. 5k,l and 6e–h). This means that for events to occur (i.e., for the FWI to exceed the 99th percentile threshold), maximum temperature anomalies compared to the contemporaneous 35-yr mean are only slightly warmer. However, precipitation anomalies fall further below the mean throughout the Pacific Coast region in the future periods (up to 10% lower in some locations), with larger decreases occurring in the eastern part of the region (Figs. 5e,f and 6a–d). Some of these larger negative anomalies are occurring where mean precipitation is increasing in future climates (supplemental Fig. 3), causing extreme fire weather events to be relatively much drier than the moving mean. Events on the coast are also becoming more humid, while interior events become less humid, compared to the moving mean, and events in the north are less windy, while events in the south see small but

robust increases in wind speed compared to the moving mean (Figs. 5w,x and 6i–p).

Events identified using a fixed threshold in the Four Corners region also show large future increases in maximum temperature of up to 4 K (Fig. 7c). The largest increases in maximum temperature occur in Utah and Colorado, coinciding with decreases in relative humidity and increases in wind speed (Figs. 7c,e,g and 8e,f,i,j,m,n). Events in the Rocky Mountains have the largest negative anomalies in precipitation and relative humidity in the historic period, and these variables continue to decrease in the future periods by 0.15 mm day^{-1} and 2%, respectively—however, this area shows relatively few event days per year in the historic period as well as smaller increases in the number of event days per year compared to other regions (Figs. 3d–f and 7a,e). Therefore, while these events are becoming drier, lower precipitation and relative humidity are not necessarily driving increases in extreme fire weather events over this region. We also note that




FIG. 8. Time series of (a)–(d) precipitation, (e)–(h) maximum temperature, (i)–(l) relative humidity, and (m)–(p) wind speed anomalies during extreme fire weather events in JJASO in (a),(e),(i),(m) UT, (b),(f),(j),(n) CO, (c),(g),(k),(o) AZ, and (d),(h),(l),(p) NM using the fixed mean and threshold (coral) and the moving mean and threshold (teal).

actual vapor pressure and VPD anomalies increase throughout the whole region during extreme fire weather events, which is reflected in changes in relative humidity in Colorado and Utah, but oppose the relative humidity trends in New Mexico and Arizona (supplemental Fig. 5). As in the Pacific Coast, events are generally less windy, with about 0.5 m s^{-1} decreases in wind speed over Utah and Colorado (Figs. 7g and 8m,n). Patterns similar to those in the Pacific Coast region also emerge for events identified using a moving threshold in the Four Corners region. Events are much drier, especially in the center of the region where mean precipitation is increasing (supplemental Fig. 3), and Utah and western Colorado show large decreases in relative humidity (Figs. 7f and 8i,j). Events have greater wind speeds in New Mexico and Arizona, but lower wind speeds in the northern part of the Four Corners region (Figs. 7h and 8m–p). Additionally, increases in maximum temperature are small and not significant for many parts of the region (Figs. 7d and 8a–d).

d. Isolating the impact of individual fire weather variables

While the previous results show how the underlying conditions of extreme fire weather events are projected to change, it

is still unclear how changes in the mean and variability of each FWI input variable impact event characteristics. To investigate this, we use quantile and mean mapping approaches (see methods, Fig. 2 and supplemental Fig. 2) to isolate the impact of total changes, as well as changes in the mean and variability of maximum temperature, precipitation, relative humidity, and wind speed on the frequency, duration, area, and intensity of extreme fire weather events (identified using the fixed threshold) in the future periods for the Pacific Coast and Four Corners regions.

In the Pacific Coast region, mean increases in maximum temperature lead to the largest increases in extreme fire weather event frequency (up to ~ 1 event per year in some locations), number of event days (~ 10), area ($\sim 10\,000\,000 \text{ km}^2$), and maximum intensity (~ 3 FWI units) in the late future period, while changes in the variability of maximum temperature have minimal impacts on these event characteristics (Figs. 9e,f and supplemental Fig. 6). For all events in the Pacific Coast, mean increases in temperature lead to the largest increases in number of events, number of events days, and area compared to any changes in other variables (Figs. 9e and 10b–d and supplemental Figs. 3 and 6).




FIG. 9. The effect of changes in the (a),(d),(g),(j) distribution, (b),(e),(h),(k) mean, and (c),(f),(i),(l) variability of (a)–(c) precipitation, (d)–(f) maximum temperature, (g)–(i) relative humidity, and (j)–(l) wind speed on changes (from the historic period, 1980–2014) in the number of event days per year for the Pacific Coast region for JJASO in the late future period (2050–82) (see supplemental Fig. 2 for details of the effect calculation for each FWI variable). These composites are created by averaging event characteristics for all events for each ensemble member for each period at each grid point. Then, the gridpoint-specific values of the historic period are subtracted from the late future period for each ensemble member, and the difference values are averaged over all ensemble members. Stippling indicates that the difference from the historic impacts is not significantly different ($p < 0.05$). Results for event frequency, area, and intensity can be found in supplemental Fig. 6.

Compared to changes in mean maximum temperature, changes in mean relative humidity cause slightly smaller increases in the number of event days and area of extreme fire weather events in the late future period over the Pacific Coast region (Figs. 9h and 10b,c and supplemental Fig. 6). While changes in mean relative humidity do lead to increases in the number of events and maximum FWI intensity in eastern Oregon and Washington, they also lead to decreases in the coastal areas and in much of California (up to -0.8 events per year and -4 FWI units in maximum FWI; supplemental Fig. 6). Unlike for maximum temperature, we find that the changes in variability (obtained as the difference between the effects of changes in distribution and mean; see methods and supplemental Fig. 2)

of relative humidity also lead to substantial and significant changes in event characteristics in much of the Pacific Coast and tend to amplify the total impact of relative humidity changes on extreme fire weather events (Figs. 9i and 10, and supplemental Fig. 6). This is especially evident for event duration and area, where changes in the mean and variability of relative humidity lead to increases of 0.25 and 0.1 event days per year and 80 000 and 20 000 km^2 in average event area, respectively, in the late future period (Figs. 10b,c). By further isolating the role of vapor pressure changes, we find that much of the increases in event days, area, and intensity are driven by the impact of rising temperatures on relative humidity (supplemental Fig. 8). Changes in vapor pressure alone lead to large decreases in these event characteristics, further highlighting the role of increasing temperatures on extreme fire weather events.

We note that changes in mean relative humidity leads to a spatially averaged reduction in number of events per year (Fig. 10a). These spatially averaged decreases are driven by large decreases on the Washington (WA) coast and broad areas of decreases in California (supplemental Fig. 6). However, when considering the role of all variables combined, these decreases are opposed by larger increases due to changes in precipitation, maximum temperature, and wind speed, especially in Southern California (CA) and WA (supplemental Fig. 6). The change in the mean of wind speed generally drives small decreases in event duration and area over the region but causes increases in the event frequency and maximum intensity of events in Southern California (Fig. 10 and supplemental Fig. 6). We hypothesize that higher mean wind speeds in Southern California are leading to higher values of FWI when extreme fire weather events do occur but may also lead to the attrition and separation of events (supplemental Fig. 3). The impact of the changes in the variability of wind speed on event characteristics is relatively small in the region except for the maximum intensity of events in Northern California where there is an increase of about 2 FWI units (Figs. 9i and 10 and supplemental Fig. 6).

In the Four Corners regions, the impacts of changes in the mean and variability of the FWI variables are somewhat similar to those in the Pacific Coast region. Mean changes in maximum temperature and relative humidity lead to large increases in event days per year (up to 15 and 8 days per year, respectively) and area (up to 10 000 000 and 8 000 000 km^2 , respectively) in the late future period (Figs. 11e,h and supplemental Fig. 7). Similar to the Pacific Coast region, the changes in mean relative humidity in the Four Corners region are likely largely driven by the impact of increasing temperatures on relative humidity, given that changes in the mean of vapor pressure lead to substantial decreases in event days and area (supplemental Fig. 9). The impact of changes in the variability of maximum temperature is relatively small. However, changes in the variability of relative humidity lead to substantial and robust increases in event days per year and event area (Fig. 11i and supplemental Fig. 7)—approximately equal to the increases due to changes in the mean of relative humidity when looking across all events and ensemble members (Figs. 12b,c). Changes in the mean and variability of relative humidity also lead to spatially varied impacts on event intensity in the Four Corners region (supplemental Fig. 6). For example, in the center of the region, there are small




FIG. 10. The difference (from the historic period, 1980–2014) in the impact of changes in the full distribution (Dist.), mean, and variability (Var.) of precipitation, relative humidity, maximum temperature, and wind speed on (a) events per year, (b) events days per year, (c) regional event area, and (d) maximum regional event intensity FWI using the fixed threshold for the early (2015–50, coral) and late future periods (2050–82, red) for the Pacific Coast region in JJASO (see supplemental Fig. 2 for details of the effect calculation for each FWI variable). The mean across all events for each ensemble member is first calculated, and we show the median across all ensemble members. The error bar represents 25th–75th percentile of the ensemble spread.




FIG. 11. The effect of changes in the (a),(d),(g),(j) distribution, (b),(e),(h),(k) mean, and (c),(f),(i),(l) variability of (a)–(c) precipitation, (d)–(f) maximum temperature, (g)–(i) relative humidity, and (j)–(l) wind speed on changes (from the historic period, 1980–2014) in the number of event days per for the Four Corners region for JJASO in the late future period (2050–82) (see supplemental Fig. 2 for details of the effect calculation for each FWI variable). These composites are created by averaging event characteristics for all events for each ensemble member for each period at each grid point. Then, the gridpoint-specific values of the historic period are subtracted from the late future period for each ensemble member, and the difference values are averaged over all ensemble members. Stippling indicates that the difference from the historic impacts is not significantly different ($p < 0.05$). Results for event frequency, area, and intensity can be found in supplemental Fig. 7.

and not statistically significant changes in maximum intensity due to changes in the mean of relative humidity but large and significant increases (approximately 6 FWI units, p value ≤ 0.05) due to changes in the variability. However, overall, changes in the mean of relative humidity lead to slight and nonrobust increases in maximum event intensity, but changes in the variability lead to large decreases (-0.5 FWI units in the late future period; Fig. 12d). Like the Pacific Coast, changes in the mean of wind speed also lead to overall decreases in the number of event days per year (-4 event days per year) and event area (-4000000 km²) but are generally overwhelmed by the increases due to changes in other variables (Fig. 11k and supplemental Fig. 9). However, in western Colorado, the large decreases in event intensity due to mean changes in wind speed (up to -10 FWI units) are also reflected in the changes due to all variables (up to -2 FWI units; supplemental Fig. 7).

In both the Pacific Coast and Four Corners regions, changes in the mean of maximum temperature and in the mean and variability of relative humidity have the largest influences on extreme fire weather event frequency, duration, area, and intensity. On the other hand, the impact of changes in wind speed

is generally small, not significant, and varied, and changes in precipitation also play a relatively small role in the changes of extreme fire weather event characteristics for both early and late future periods.

4. Discussion

Our study employs the CESM2 Large Ensemble to investigate projections of extreme fire weather event characteristics for current and emerging fire-prone regions, specifically, the Pacific Coast and Four Corners regions. Using a fixed threshold, we find that the frequency, duration, and area of extreme fire weather events are projected to increase under a high-emission scenario and projected to extend well-beyond regional boundaries. By the end of the twenty-first century, events in the Pacific Region are expected to reach north toward Canada and east toward the Midwest, and events in the Four Corners regions are expected to reach northeast toward the Midwest and west and northwest toward Idaho, Nevada, and Southern California. These increases in extreme fire weather event duration, area, and intensity are fueled by increasing maximum temperature means and decreasing relative humidity means, as well as changes in the variability of relative humidity that are not reflected in the mean changes. When using a moving threshold, event frequency, duration, and area are generally unchanging in the future periods, but some expansion of spatiotemporally connected events toward similar regions as the fixed thresholds is evident, reflecting changes that are not explained by the long-term, mean climate trend. Additionally, extreme fire weather events using a moving threshold are only slightly warmer than the contemporaneous mean maximum temperature but occur under much drier conditions, reflected in the underlying anomalously low precipitation and relative humidity. These changes may reflect the increases in mean precipitation throughout the future period that result in larger anomalies during extreme fire weather events.

We find that the choice of the baseline or threshold period used to investigate extreme fire weather events is highly important for making inferences about future events. In our case, we used two contrasting baselines: a fixed threshold that relies on the distribution of the FWI in the historical period only and a moving threshold that accounts for an evolving FWI distribution. The relevance of a fixed versus moving baseline for understanding the risk of extreme fire weather events in a particular region depends on the time scales at which communities and ecosystems adapt to changes in mean climate versus climate variability. The moving threshold also allows insight into the drivers of future extreme fire weather events beyond the thermodynamically imposed shift in the FWI distribution. Similar considerations apply to other types of extremes, for example, marine (Deser et al. 2024; Amaya et al. 2023; Capotondi et al. 2024; Smith et al. 2025) and terrestrial heat waves (Skinner et al. 2025; Vogel et al. 2020). As far as we know, this is the first study to explicitly compare the impacts of projected changes in mean climate versus climate variability on the characteristics of extreme fire weather events and to quantify the underlying meteorological drivers of such changes.




FIG. 12. The difference (from the historic period, 1980–2014) in the impact of changes in the full distribution (Dist.), mean, and variability (Var.) of precipitation, relative humidity, maximum temperature, and wind speed on (a) events per year, (b) events days per year, (c) regional event area, and (d) maximum regional event intensity FWI using the fixed threshold for the early (2015–50, coral) and late future periods (2050–82, red) for the Four Corners region in JJASO (see supplemental Fig. 2 for details of the effect calculation for each FWI variable). The mean across all events for each ensemble member is first calculated, and we show the median across all ensemble members. The error bar represents 25th–75th percentile of the ensemble spread.

By using a moving threshold, we can bring to light the changes in event characteristics that may be due to changes in spatial or temporal variability of the FWI distribution that are not obvious when using a fixed threshold. In other words, we are able to isolate the changes in extreme fire weather events that are not due to the thermodynamically driven trends in the FWI and that may act on relatively shorter time scales. We find some changes in extreme fire weather event characteristics when using a moving threshold, namely, the underlying meteorological conditions that lead to extreme fire weather events, as well as the spatial connectivity of events in both the Pacific Coast and Four Corners regions. For example, we show that when considering a moving window to define the threshold and mean climate, extreme fire weather events are much drier but not substantially warmer, showing that on short time scales, anomalously low precipitation is more impactful than anomalously high temperatures.

To our knowledge, our study is the first to employ a quantile mapping method to isolate the changes in the mean and variability of the FWI input variables to understand how they individually impact extreme fire weather events. Moreover, we have further isolated the role of changes in vapor pressure on extreme fire weather events through its impact on changes in relative humidity. The use of a single model introduces physics-based uncertainties and uncertainties due to climate sensitivities. However, based on the analysis in this manuscript (supplemental Fig. 1) and in previous studies, we are confident that the model captures the FWI reasonably well and projects the FWI in the future similarly to other CMIP-style models (Touma et al. 2022; Gallo et al. 2023). Notably, the CESM2-LE allows for the robust estimate of anthropogenically forced changes in the mean and variability of the individual variables due to the availability of 100 simulations that differ only by internal variability. While our findings show the large impact of increases in the mean of maximum temperatures and decreases in the mean of relative humidity that have been shown in previous studies (Abatzoglou et al. 2019; Jain et al. 2022; Touma et al. 2021), they also show that changes in the variability of relative humidity have a similarly large impact on extreme fire weather event frequency, duration, area, and intensity. In some areas of our study region, the impact of changes in the variability of relative humidity is equal to that of changes in the mean, doubling the total impact of relative humidity, which is a substantially larger estimate than that of previous studies (e.g., Zhuang et al. 2021). Our findings stress the need to consider changes in the variability as well as the mean of climate variables when understanding climate change-induced changes in different characteristics of extreme climate events and, therefore, to employ large ensembles for assessing projections of extreme events.

By using a 35-yr moving window that includes all daily values in our quantile-mapping method, we do not differentiate between different time scales of variability—daily-to-multiannual time scales are considered together. Previous studies have shown that El Niño–Southern Oscillation (ENSO) and the Pacific decadal oscillation (PDO) influence the variability of wildfire occurrence over the preindustrial and historical period in the western United States and that the Atlantic meridional oscillation (AMO) can also modulate ENSO's and PDO's

influences (Kitzberger et al. 2007). Studies have also shown that ENSO's influence on wildfire danger in the western United States is expected to change in future years, as well as the influence on different wildfire-related variables (Fasullo et al. 2018).

5. Conclusions

Our study is one of many that shows that western U.S. wildfire risk is projected to increase over the next century due to increased frequency, persistence, and areal coverage of extreme fire weather conditions and that communities need to plan how to become resilient to these changes. By employing the moving threshold and understanding the role of variability and mean changes in our study, we also begin to shed light on how adaptation measures should be shaped. For instance, our study showed that extreme fire weather events are not only projected to increase in frequency but also likely to spread over larger areas and become more spatially connected over the coming decades, highlighting the need to prepare for more widespread hazardous conditions. We also found that these changes are not only thermodynamically driven and that robust and significant changes also occur beyond those that are driven by changes in the mean climate. This finding provides insight into the drivers and predictability of extreme fire weather events on subseasonal time scales that are masked when using a fixed threshold (Smith et al. 2025). Our study also shows that by using a large ensemble, we capture the impact of forced changes in the variability of climate variables on extreme fire weather events.

While the use of regionally downscaled (statistically and dynamically) climate projections is important for adaptation policy, they rarely include more than a few realizations and, therefore, could hide the impacts of the forced changes in climate variability [see Deser et al. (2020) for further reading]. One way to overcome this issue is to statistically downscale global ESM large ensembles that are run at relatively coarse resolution or select a few “representative” ensemble members that could capture the forced changes in variability and use those to dynamically downscale climate projections (see, e.g., Huang and Swain 2022). While both pathways will lead to their own uncertainties, we stress the need to capture the impact of anthropogenic forcings on both the variability and mean of our climate system to quantify future risks of extreme fire weather events among other impacts.

Acknowledgments. We thank five anonymous reviewers and the editor, Dr. Andrew Hoell, for insightful and constructive feedback. This material is based upon work supported by the NSF National Center for Atmospheric Research, which is a major facility sponsored by the U.S. National Science Foundation under Cooperative Agreement 1852977. We would like to acknowledge computing support from the Casper system (<https://ncar.pub/casper>) provided by the NSF National Center for Atmospheric Research (NCAR), sponsored by the National Science Foundation. We also acknowledge the CESM2 Large Ensemble Community Project and supercomputing resources provided by the IBS Center for Climate Physics in South Korea.

Data availability statement. The CESM2-LE model output is available through <https://www.cesm.ucar.edu/projects/community-projects/LENS2/data-sets.html>. Analysis and visualization scripts are available at https://github.com/detouma/Touma_Deser_JClim_2026.

REFERENCES

- Abatzoglou, J., M. Jones, C. Kolden, A. Cullen, M. Sadegh, and E. Williams, 2025: Climate change has increased the odds of extreme regional forest fire years globally. *EGU General Assembly 2025*, Vienna, Austria, European Geosciences Union, Abstract EGU25-2594, <https://doi.org/10.5194/egusphere-egu25-2594>.
- Abatzoglou, J. T., 2013: Development of gridded surface meteorological data for ecological applications and modelling. *Int. J. Climatol.*, **33**, 121–131, <https://doi.org/10.1002/joc.3413>.
- , and A. P. Williams, 2016: Impact of anthropogenic climate change on wildfire across western US forests. *Proc. Natl. Acad. Sci. USA*, **113**, 11 770–11 775, <https://doi.org/10.1073/pnas.1607171113>.
- , —, and R. Barbero, 2019: Global emergence of anthropogenic climate change in fire weather indices. *Geophys. Res. Lett.*, **46**, 326–336, <https://doi.org/10.1029/2018GL080959>.
- , C. S. Juang, A. P. Williams, C. A. Kolden, and A. L. Westerling, 2021: Increasing synchronous fire danger in forests of the western United States. *Geophys. Res. Lett.*, **48**, e2020GL091377, <https://doi.org/10.1029/2020GL091377>.
- Amaya, D. J., and Coauthors, 2023: Marine heatwaves need clear definitions so coastal communities can adapt. *Nature*, **616**, 29–32, <https://doi.org/10.1038/d41586-023-00924-2>.
- Andela, N., and Coauthors, 2017: A human-driven decline in global burned area. *Science*, **356**, 1356–1362, <https://doi.org/10.1126/science.aal4108>.
- Bui, H. X., Y.-X. Li, and D. Domménget, 2024: Controlling factors of wildfires in Australia and their changes under global warming. *Environ. Res. Lett.*, **19**, 094030, <https://doi.org/10.1088/1748-9326/ad69a9>.
- Burke, M., A. Driscoll, S. Heft-Neal, J. Xue, J. Burney, and M. Wara, 2021: The changing risk and burden of wildfire in the United States. *Proc. Natl. Acad. Sci. USA*, **118**, e2011048118, <https://doi.org/10.1073/pnas.2011048118>.
- Capotondi, A., and Coauthors, 2024: A global overview of marine heatwaves in a changing climate. *Commun. Earth Environ.*, **5**, 701, <https://doi.org/10.1038/s43247-024-01806-9>.
- Collar, N. M., S. Saxe, B. A. Ebel, K. S. Boden, A. J. Rust, and T. S. Hogue, 2022: Linking fire-induced evapotranspiration shifts to streamflow magnitude and timing in the western United States. *J. Hydrol.*, **612**, 128242, <https://doi.org/10.1016/j.jhydrol.2022.128242>.
- Cook, B. I., and R. Seager, 2013: The response of the North American monsoon to increased greenhouse gas forcing. *J. Geophys. Res. Atmos.*, **118**, 1690–1699, <https://doi.org/10.1002/jgrd.50111>.
- DeRepentigny, P., and Coauthors, 2022: Enhanced simulated early 21st century Arctic sea ice loss due to CMIP6 biomass burning emissions. *Sci. Adv.*, **8**, eabo2405, <https://doi.org/10.1126/sciadv.abo2405>.
- Deser, C., and Coauthors, 2020: Insights from Earth system model initial-condition large ensembles and future prospects. *Nat. Climate Change*, **10**, 277–286, <https://doi.org/10.1038/s41558-020-0731-2>.
- , A. S. Phillips, M. A. Alexander, D. J. Amaya, A. Capotondi, M. G. Jax, and J. D. Scott, 2024: Future changes in the intensity and duration of marine heat and cold waves: Insights from coupled model initial-condition large ensembles. *J. Climate*, **37**, 1877–1902, <https://doi.org/10.1175/JCLI-D-23-0278.1>.
- , W. M. Kim, R. C. J. Wills, I. R. Simpson, S. Yeager, G. Danabasoglu, K. Rodgers, and N. Rosenbloom, 2025: Effects of macro vs. micro initialization and ocean initial-condition memory on the evolution of ensemble spread in the CESM2 large ensemble. *Climate Dyn.*, **63**, 62, <https://doi.org/10.1007/s00382-024-07553-z>.
- Dowdy, A., G. A. Mills, K. Finkele, and W. De Groot, 2009: Australian fire weather as represented by the McArthur Forest Fire Danger Index and the Canadian Forest Fire Weather Index. CAWCR Tech. Rep. 10, 91 pp.
- Fasullo, J. T., 2020: Evaluating simulated climate patterns from the CMIP archives using satellite and reanalysis datasets using the Climate Model Assessment Tool (CMATv1). *Geosci. Model Dev.*, **13**, 3627–3642, <https://doi.org/10.5194/gmd-13-3627-2020>.
- , B. L. Otto-Bliesner, and S. Stevenson, 2018: ENSO's changing influence on temperature, precipitation, and wildfire in a warming climate. *Geophys. Res. Lett.*, **45**, 9216–9225, <https://doi.org/10.1029/2018GL079022>.
- Gallo, C., J. M. Eden, B. Dieppois, I. Drobyshev, P. Z. Fulé, J. San-Miguel-Ayanz, and M. Blackett, 2023: Evaluation of CMIP6 model performances in simulating fire weather spatiotemporal variability on global and regional scales. *Geosci. Model Dev.*, **16**, 3103–3122, <https://doi.org/10.5194/gmd-16-3103-2023>.
- Goss, M., D. L. Swain, J. T. Abatzoglou, A. Sarhadi, C. A. Kolden, A. P. Williams, and N. S. Diffenbaugh, 2020: Climate change is increasing the likelihood of extreme autumn wildfire conditions across California. *Environ. Res. Lett.*, **15**, 094016, <https://doi.org/10.1088/1748-9326/ab83a7>.
- Hagmann, R. K., and Coauthors, 2021: Evidence for widespread changes in the structure, composition, and fire regimes of western North American forests. *Ecol. Appl.*, **31**, e02431, <https://doi.org/10.1002/eap.2431>.
- Hoell, A., and Coauthors, 2022: Record low North American monsoon rainfall in 2020 reignites drought over the American southwest. *Bull. Amer. Meteor. Soc.*, **103** (3), S26–S32, <https://doi.org/10.1175/BAMS-D-21-0129.1>.
- Huang, X., and D. L. Swain, 2022: Climate change is increasing the risk of a California megaflood. *Sci. Adv.*, **8**, eabq0995, <https://doi.org/10.1126/sciadv.abq0995>.
- Jain, P., D. Castellanos-Acuna, S. C. P. Coogan, J. T. Abatzoglou, and M. D. Flannigan, 2022: Observed increases in extreme fire weather driven by atmospheric humidity and temperature. *Nat. Climate Change*, **12**, 63–70, <https://doi.org/10.1038/s41558-021-01224-1>.
- , and Coauthors, 2024: Drivers and impacts of the record-breaking 2023 wildfire season in Canada. *Nat. Commun.*, **15**, 6764, <https://doi.org/10.1038/s41467-024-51154-7>.
- Jones, M. W., and Coauthors, 2022: Global and regional trends and drivers of fire under climate change. *Rev. Geophys.*, **60**, e2020RG000726, <https://doi.org/10.1029/2020RG000726>.
- , and Coauthors, 2024: State of wildfires 2023–2024. *Earth Syst. Sci. Data*, **16**, 3601–3685, <https://doi.org/10.5194/essd-16-3601-2024>.
- Juang, C. S., A. P. Williams, J. T. Abatzoglou, J. K. Balch, M. D. Hurteau, and M. A. Moritz, 2022: Rapid growth of large forest fires drives the exponential response of annual forest-fire area

- to aridity in the western United States. *Geophys. Res. Lett.*, **49**, e2021GL097131, <https://doi.org/10.1029/2021GL097131>.
- Kirchmeier-Young, M. C., F. W. Zwiers, N. P. Gillett, and A. J. Cannon, 2017: Attributing extreme fire risk in western Canada to human emissions. *Climate Change*, **144**, 365–379, <https://doi.org/10.1007/s10584-017-2030-0>.
- , and Coauthors, 2024: Human driven climate change increased the likelihood of the 2023 record area burned in Canada. *npj Climate Atmos. Sci.*, **7**, 316, <https://doi.org/10.1038/s41612-024-00841-9>.
- Kitzberger, T., P. M. Brown, E. K. Heyerdahl, T. W. Swetnam, and T. T. Veblen, 2007: Contingent Pacific–Atlantic Ocean influence on multicentury wildfire synchrony over western North America. *Proc. Natl. Acad. Sci. USA*, **104**, 543–548, <https://doi.org/10.1073/pnas.0606078104>.
- Lawrence, A. J., C. Matuch, J. J. Hancock, A. L. Rypel, and L. A. Eliassen, 2022: Potential local extirpation of an imperiled freshwater mussel population from wildfire runoff. *West. North Amer. Nat.*, **82**, 695–703, <https://doi.org/10.3398/064.082.0405>.
- Oakley, N. S., 2021: A warming climate adds complexity to post-fire hydrologic hazard planning. *Earth's Future*, **9**, e2021EF002149, <https://doi.org/10.1029/2021EF002149>.
- Raymond, C., and Coauthors, 2020: Understanding and managing connected extreme events. *Nat. Climate Change*, **10**, 611–621, <https://doi.org/10.1038/s41558-020-0790-4>.
- Rodgers, K. B., and Coauthors, 2021: Ubiquity of human-induced changes in climate variability. *Earth Syst. Dyn.*, **12**, 1393–1411, <https://doi.org/10.5194/esd-12-1393-2021>.
- Sharma, A. R., P. Jain, J. T. Abatzoglou, and M. Flannigan, 2022: Persistent positive anomalies in geopotential heights promote wildfires in western North America. *J. Climate*, **35**, 6469–6486, <https://doi.org/10.1175/JCLI-D-21-0926.1>.
- Simpson, I. R., K. A. McKinnon, D. Kennedy, D. M. Lawrence, F. Lehner, and R. Seager, 2024: Observed humidity trends in dry regions contradict climate models. *Proc. Natl. Acad. Sci. USA*, **121**, e2302480120, <https://doi.org/10.1073/pnas.2302480120>.
- Skinner, C. B., D. Touma, M. Barlow, D. Singh, and T. King, 2025: The spatial extent of heat waves has changed over the past four decades. *Commun. Earth Environ.*, **6**, 662, <https://doi.org/10.1038/s43247-025-02661-y>.
- Smith, K. E., and Coauthors, 2025: Baseline matters: Challenges and implications of different marine heatwave baselines. *Prog. Oceanogr.*, **231**, 103404, <https://doi.org/10.1016/j.pocean.2024.103404>.
- Touma, D., S. Stevenson, F. Lehner, and S. Coats, 2021: Human-driven greenhouse gas and aerosol emissions cause distinct regional impacts on extreme fire weather. *Nat. Commun.*, **12**, 212, <https://doi.org/10.1038/s41467-020-20570-w>.
- , —, D. L. Swain, D. Singh, D. A. Kalashnikov, and X. Huang, 2022: Climate change increases risk of extreme rainfall following wildfire in the western United States. *Sci. Adv.*, **8**, eabm0320, <https://doi.org/10.1126/sciadv.abm0320>.
- , J. W. Hurrell, M. R. Tye, and K. Dagon, 2023: The impact of stratospheric aerosol injection on extreme fire weather risk. *Earth's Future*, **11**, e2023EF003626, <https://doi.org/10.1029/2023EF003626>.
- van Marle, M. J. E., and Coauthors, 2017: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015). *Geosci. Model Dev.*, **10**, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>.
- Van Wagner, C. E., 1987: Development and structure of the Canadian Forest Fire Weather Index System. Minister of Supply and Services Canada Forestry Tech. Rep. 35, 35 pp.
- Virtanen, P., and Coauthors, 2020: SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nat. Methods*, **17**, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>.
- Vogel, M. M., J. Zscheischler, E. M. Fischer, and S. I. Seneviratne, 2020: Development of future heatwaves for different hazard thresholds. *J. Geophys. Res. Atmos.*, **125**, e2019JD032070, <https://doi.org/10.1029/2019JD032070>.
- Williams, A. P., and Coauthors, 2022: Growing impact of wildfire on western US water supply. *Proc. Natl. Acad. Sci. USA*, **119**, e2114069119, <https://doi.org/10.1073/pnas.2114069119>.
- Zhuang, Y., R. Fu, B. D. Santer, R. E. Dickinson, and A. Hall, 2021: Quantifying contributions of natural variability and anthropogenic forcings on increased fire weather risk over the western United States. *Proc. Natl. Acad. Sci. USA*, **118**, e2111875118, <https://doi.org/10.1073/pnas.2111875118>.
- Zscheischler, J., and Coauthors, 2020: A typology of compound weather and climate events. *Nat. Rev. Earth Environ.*, **1**, 333–347, <https://doi.org/10.1038/s43017-020-0060-z>.